

## ENVELOPES FOR SETS AND FUNCTIONS: REGULARIZATION AND GENERALIZED CONJUGACY

A. CABOT, A. JOURANI AND L. THIBAUT

*Abstract.* Let  $X$  be a vector space and let  $\varphi : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  be an extended real-valued function. For every function  $f : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ , let us define the  $\varphi$ -envelope of  $f$  by

$$f^\varphi(x) = \sup_{y \in X} \varphi(x - y) \dot{-} f(y),$$

where  $\dot{-}$  denotes the lower subtraction in  $\mathbb{R} \cup \{-\infty, +\infty\}$ . The main purpose of this paper is to study in great detail the properties of the important generalized conjugation map  $f \mapsto f^\varphi$ . When the function  $\varphi$  is closed and convex,  $\varphi$ -envelopes can be expressed as Legendre–Fenchel conjugates. By particularizing with  $\varphi = (1/p\lambda)\|\cdot\|^p$ , for  $\lambda > 0$  and  $p \geq 1$ , this allows us to derive new expressions of the Klee envelopes with index  $\lambda$  and power  $p$ . Links between  $\varphi$ -envelopes and Legendre–Fenchel conjugates are also explored when  $-\varphi$  is closed and convex. The case of Moreau envelopes is examined as a particular case. In addition to the  $\varphi$ -envelopes of functions, a parallel notion of envelope is introduced for subsets of  $X$ . Given subsets  $\Lambda, C \subset X$ , we define the  $\Lambda$ -envelope of  $C$  as  $C^\Lambda = \bigcap_{x \in C} (x + \Lambda)$ . Connections between the transform  $C \mapsto C^\Lambda$  and the aforesaid  $\varphi$ -conjugation are investigated.

**§1. Introduction.** Given two topological vector spaces  $X, Y$  and a function  $c : X \times Y \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ , extending the Legendre–Fenchel conjugacy, Moreau [22, Ch. 14, §3] defined, for any function  $g : Y \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  its  $c$ -conjugate as the function  $g^c : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ ,

$$g^c(x) := \sup_{y \in Y} (c(x, y) \dot{-} g(y)) \quad \text{for all } x \in X;$$

see §2 for the (extended) lower subtraction  $\dot{-}$ . We refer to [4, 6, 7, 9, 17, 22, 31, 37] and the references therein for various duality results in such a context and for several applications. Given a function  $\varphi : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  we will focus on the case  $c(x, y) := \varphi(x - y)$  and  $Y = X$ . Otherwise stated, for a function  $f : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  we will be interested in the function  $f^\varphi$  that we call the  $\varphi$ -envelope of  $f$ , defined by

$$f^\varphi(x) := \sup_{y \in X} (\varphi(x - y) \dot{-} f(y)) \quad \text{for all } x \in X.$$

Our first aim in this paper is to study in great detail the structure of the transform  $f \mapsto f^\varphi$  and provide various properties of  $\varphi$ -envelopes.

---

Received 8 April 2016.  
 MSC (2010): 52A41, 49J53, 41A65 (primary).

On the other hand, considering the class  $\mathcal{B}_X$  of closed balls of a Banach space  $X$ , Mazur [19] studied some Banach spaces  $X$  for which every closed bounded convex subset is the intersection of some subclass of  $\mathcal{B}_X$ ; we refer to [10] for a rich survey on the subject. Any such Banach space is actually called in the literature a Banach space with the Mazur intersection property. In his 1983 paper [36] Vial defined strongly convex sets of a normed space as convex sets which are intersections of closed balls with a common radius; sets which are intersections, for a fixed real  $r > 0$ , of closed balls with radius equal to  $r$  are called  $r$ -strongly convex sets in [36]. This class of convex sets is thoroughly studied by Polovinkin [25] (see also [26] and the references therein). Denoting by  $\mathbb{B}_X$  the closed unit ball of  $X$  centered at zero, any  $r$ -strongly convex set can be represented in the form

$$\bigcap_{x \in S} (x + r\mathbb{B}_X) \quad \text{with some subset } S \subset X.$$

So, given a subset  $\Lambda$  of the space  $X$ , our second aim in the paper is to analyze properties of the transform which assigns to each subset  $C$  of  $X$  the set

$$C^\Lambda := \bigcap_{x \in C} (x + \Lambda).$$

We will also provide the connections between the latter transform and the aforestated transform related to  $\varphi$ -envelopes.

In §2 we recall the lower and upper additions (respectively subtractions), and we also recall various concepts and results in convex analysis which will be needed in the paper. Section 3 offers a large list of general properties of  $\varphi$ -envelopes. Section 4 establishes the connections between  $\varphi$ -envelopes and the aforementioned transform  $C \mapsto C^\Lambda$ ; many properties of sets which can be represented in this form are also provided. In §5 we examine the question whether  $\psi = \varphi(\cdot - a) - \alpha$  (for some  $a \in X$  and  $\alpha \in \mathbb{R}$ ) whenever  $\psi$  is a  $\varphi$ -envelope and  $\varphi$  is a  $\psi$ -envelope. A counter-example is constructed and various sufficient conditions are given. The analogous question is also investigated with sets instead of functions. Section 6 considers additional properties in the case when the function  $\varphi$  is either superadditive or subadditive. In §7, assuming that  $\varphi$  is convex and lower semicontinuous, we provide several links between the  $\varphi$ -envelope of a function and Legendre–Fenchel conjugates of other functions related to  $f$ . Taking  $\varphi$  as a power of the norm, we also provide various results concerning the Klee envelope  $\kappa_{\lambda,p}f$  (with index  $\lambda$  and power  $p$ ) of a function  $f$ , where

$$\kappa_{\lambda,p}f(x) := \sup_{y \in X} \left( \frac{1}{p\lambda} \|x - y\|^p - f(y) \right) \quad \text{for all } x \in X.$$

Finally in §8, assuming that  $-\varphi$  is convex and lower semicontinuous, we continue to explore the links between  $\varphi$ -envelopes and Legendre–Fenchel conjugates. By particularizing with  $\varphi = -(1/p\lambda)\|\cdot\|^p$ , for  $\lambda > 0$  and  $p \geq 1$ , we obtain several properties of Moreau envelopes with index  $\lambda$  and power  $p$ .

§2. *Preliminaries.* Following Moreau [22], we extend the usual addition on  $\mathbb{R}$  to  $\overline{\mathbb{R}} = [-\infty, +\infty]$ . We define the upper addition  $\dot{+}$  and the lower addition  $\dot{-}$  as the laws extending the usual addition via the following conventions:

$$\begin{aligned} (-\infty) \dot{+} (+\infty) &= (+\infty) \dot{+} (-\infty) = +\infty, \\ (-\infty) \dot{-} (+\infty) &= (+\infty) \dot{-} (-\infty) = -\infty. \end{aligned}$$

This leads to the introduction of the upper subtraction  $\dot{-}$  and the lower subtraction  $\dot{+}$ , respectively defined by

$$s \dot{-} t = s \dot{+} (-t) \quad \text{and} \quad s \dot{+} t = s \dot{-} (-t) \quad \text{for all } s, t \in \overline{\mathbb{R}}.$$

Let  $X$  be a vector space; all vector spaces will be real vector spaces. Given two extended real-valued functions  $f, g : X \rightarrow \overline{\mathbb{R}}$ , the (Moreau) *inf-convolution* (also called *infimal convolution*) of  $f$  and  $g$  is defined as follows: for every  $x \in X$ ,

$$\begin{aligned} (f \nabla g)(x) &= \inf_{y+z=x} [f(y) \dot{+} g(z)] \\ &= \inf_{y \in X} [f(y) \dot{+} g(x - y)] \\ &= \inf_{z \in X} [f(x - z) \dot{+} g(z)]. \end{aligned}$$

In a symmetric way, the (Moreau) *sup-convolution* (or *supremal convolution*) of  $f$  and  $g$  is defined by

$$\begin{aligned} (f \Delta g)(x) &= \sup_{y+z=x} [f(y) \dot{-} g(z)] \\ &= \sup_{y \in X} [f(y) \dot{-} g(x - y)] \\ &= \sup_{z \in X} [f(x - z) \dot{-} g(z)]. \end{aligned}$$

For the function  $f$  as above, the set  $\text{dom } f = \{x \in X, f(x) < +\infty\}$  is called the *effective domain* of  $f$ . We call  $f$  a proper function if  $f(x) < +\infty$  for at least one  $x \in X$ , and  $f(x) > -\infty$  for all  $x \in X$ , or in other words, if  $\text{dom } f$  is a non-empty set on which  $f$  is finite. The function which is constantly equal to  $+\infty$  (respectively  $-\infty$ ) on  $X$  is denoted by  $\omega_X$  (respectively  $-\omega_X$ ).

Now assume that  $X$  is a locally convex space; all such spaces in the paper will be Hausdorff. We will denote by  $X^*$  the topological dual of  $X$ . Then, following again [22], we set

$$\Gamma(X) := \{f : X \rightarrow \overline{\mathbb{R}}, f \text{ is a pointwise supremum of a family of continuous affine functions with slopes in } X^*\}$$

and

$$\Gamma(X^*) := \{g : X^* \rightarrow \overline{\mathbb{R}}, g \text{ is a pointwise supremum of a family of continuous affine functions with slopes in } X\}.$$

We denote by  $\Gamma_0(X)$  the set of  $f \in \Gamma(X)$  which differ from  $\omega_X$  and  $-\omega_X$ . In the same way,  $\Gamma_0(X^*)$  is the set  $\Gamma_0(X^*) = \Gamma(X^*) \setminus \{\omega_{X^*}, -\omega_{X^*}\}$ . The classes  $\Gamma_0(X)$  and  $\Gamma_0(X^*)$  are respectively characterized by

$$\begin{aligned} \Gamma_0(X) &= \{f : X \rightarrow \overline{\mathbb{R}}, f \text{ is closed, convex and proper}\} \\ &= \{f : X \rightarrow \overline{\mathbb{R}}, f \text{ is } w(X, X^*) \text{ closed, convex and proper}\}, \end{aligned}$$

and

$$\Gamma_0(X^*) = \{g : X^* \rightarrow \overline{\mathbb{R}}, g \text{ is } w(X^*, X) \text{ closed, convex and proper}\},$$

see, for example, [1, 8, 22]. Above and in the rest of the paper,  $w(X, X^*)$  and  $w(X^*, X)$  stand for the weak topology on  $X$  and the weak star topology on  $X^*$ , respectively.

With the function  $f : X \rightarrow \overline{\mathbb{R}}$  is associated, in the duality pairing from  $X$  to  $X^*$ , its Legendre–Fenchel conjugate  $f^* : X^* \rightarrow \overline{\mathbb{R}}$  defined by

$$\text{for all } x^* \in X^*, \quad f^*(x^*) = \sup_{\xi \in X} \{\langle x^*, \xi \rangle - f(\xi)\}.$$

In the same way, throughout the paper (unless otherwise stated) the Legendre–Fenchel conjugate of a function  $g : X^* \rightarrow \overline{\mathbb{R}}$  defined on the dual space  $X^*$  will be taken in the duality pairing from  $X^*$  to  $X$ , that is,  $g^* : X \rightarrow \overline{\mathbb{R}}$  is defined on  $X$  by

$$\text{for all } x \in X, \quad g^*(x) = \sup_{\xi^* \in X^*} \{\langle \xi^*, x \rangle - g(\xi^*)\}.$$

The Legendre–Fenchel transform  $f \mapsto f^*$  (see, for example, [22]) is known to be a one-to-one mapping from  $\Gamma_0(X)$  onto  $\Gamma_0(X^*)$ . For any  $f \in \Gamma_0(X)$  one has  $f = f^{**}$  and for any  $g \in \Gamma_0(X^*)$  one has  $g = g^{**}$  (see, for example, [1, 8, 22]).

Given a set  $C \subset X$ , we denote as usual by  $\delta_C$  the indicator function of  $C$ , i.e.  $\delta_C(x) = 0$  if  $x \in C$  and  $\delta_C(x) = +\infty$  if  $x \notin C$ . The support function  $\sigma_C : X^* \rightarrow \overline{\mathbb{R}}$  of  $C$  is defined by

$$\text{for all } x^* \in X^*, \quad \sigma_C(x^*) = \sup_{\xi \in C} \langle x^*, \xi \rangle,$$

so  $\sigma_C$  coincides with the Legendre–Fenchel conjugate of  $\delta_C$ . For a non-empty cone  $K \subset X$ , the support function  $\sigma_K$  is equal to the indicator function of the polar cone  $K^\circ$  of  $K$  defined by

$$K^\circ = \{x^* \in X^*, \langle x^*, x \rangle \leq 0 \text{ for all } x \in K\}.$$

For a set  $C \subset X$ , we denote by  $\text{co}(C)$  (respectively  $\overline{\text{co}}(C)$ ) the convex hull (respectively closed convex hull) of  $C$ . The  $w(X^*, X)$ -closed convex hull of a set  $D \subset X^*$  is denoted by  $\overline{\text{co}}^{w^*}(D)$ . For a function  $f : X \rightarrow \overline{\mathbb{R}}$ , its convex hull  $\text{co}(f)$  (respectively lower semicontinuous convex hull  $\overline{\text{co}}(f)$ ) is the greatest convex (respectively lower semicontinuous convex) function less than or equal to  $f$ . The  $w(X^*, X)$ -lower semicontinuous convex hull of a function  $g : X^* \rightarrow \overline{\mathbb{R}}$  is denoted by  $\overline{\text{co}}^{w^*}(g)$ .

If  $f \in \Gamma_0(X)$  and if  $\bar{x} \in \text{dom } f$ , the *recession function*  $f^\infty$  is defined by

$$\text{for all } u \in X, \quad f^\infty(u) = \lim_{t \rightarrow +\infty} \frac{f(\bar{x} + tu) - f(\bar{x})}{t} = \sup_{t > 0} \frac{f(\bar{x} + tu) - f(\bar{x})}{t}.$$

The function  $f^\infty : X \rightarrow \mathbb{R} \cup \{+\infty\}$  does not depend on the point  $\bar{x} \in \text{dom } f$  since it is also given by

$$\text{for all } u \in X, \quad f^\infty(u) = \sup_{x \in \text{dom } f} (f(x + u) - f(x)).$$

The function  $f^\infty$  satisfies  $f^\infty \in \Gamma_0(X)$ , it is positively homogeneous and we have  $f^\infty = \sigma_{\text{dom } f^*}$ . Given a closed convex set  $C \subset X$  and  $\bar{x} \in C$ , the recession cone  $C^\infty$  is defined by

$$C^\infty = \{u \in X, \bar{x} + tu \in C \text{ for all } t \geq 0\}.$$

The set  $C^\infty$  does not depend on  $\bar{x} \in C$  and is also given by

$$C^\infty = \{u \in X, u + C \subset C\}.$$

It follows from the definition that  $C^\infty$  is a closed convex cone and we have  $\delta_{C^\infty} = (\delta_C)^\infty$ . For more details on recession analysis, see, for example, [1, 2, 13, 28].

Let us end these preliminaries with the subdifferential of convex analysis. We recall that the *subdifferential*  $\partial f(x)$  of a convex function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  at  $x \in \text{dom } f$  is the set

$$\partial f(x) = \{\xi^* \in X^*, f(y) \geq f(x) + \langle \xi^*, y - x \rangle \text{ for every } y \in X\}. \quad (1)$$

When  $x \notin \text{dom } f$ , then  $\partial f(x) = \emptyset$  by convention. The *domain* and the *range* of the operator  $\partial f : X \rightrightarrows X^*$  are respectively given by

$$\begin{aligned} \text{dom}(\partial f) &= \{x \in X, \partial f(x) \neq \emptyset\}, \\ \text{Rge}(\partial f) &= \{x^* \in X^*, \exists x \in X, x^* \in \partial f(x)\}. \end{aligned}$$

If  $f \in \Gamma_0(X)$ , the subdifferentials of  $f$  and  $f^*$  are connected through the following relation

$$x^* \in \partial f(x) \iff x \in \partial f^*(x^*), \quad (2)$$

for all  $x \in X$  and  $x^* \in X^*$ . For further details, the reader is referred to the classical textbooks on convex analysis, see, for example, [13, 28].

§3. *Definitions, general properties.* Let  $X$  be a vector space. For functions  $\varphi : X \rightarrow \overline{\mathbb{R}}$  and  $f : X \rightarrow \overline{\mathbb{R}}$ , the  $\varphi$ -*envelope* of  $f$  is defined as follows:

$$\text{for all } x \in X, \quad f^\varphi(x) = \sup_{y \in X} \{\varphi(x - y) - f(y)\} = \sup_{z \in X} \{\varphi(z) - f(x - z)\}.$$

A function  $g : X \rightarrow \overline{\mathbb{R}}$  is said to be a  $\varphi$ -envelope if there exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $g = f^\varphi$ . It is immediate to check that for every function  $f : X \rightarrow \overline{\mathbb{R}}$ ,

$$f^{-\omega_X} = -\omega_X, \quad \text{while } f^{\omega_X} = \begin{cases} \omega_X & \text{if } f \neq \omega_X, \\ -\omega_X & \text{if } f = \omega_X. \end{cases}$$

It ensues that the unique  $(-\omega_X)$ -envelope is the function  $-\omega_X$ , while the  $\omega_X$ -envelopes are  $\pm\omega_X$ . The function  $f^\varphi$  can be expressed via the inf-convolution and sup-convolution operators

$$f^\varphi = \varphi \Delta (-f) = -((-\varphi) \nabla f). \tag{3}$$

The roles played by  $f$  and  $\varphi$  in the definition of  $f^\varphi$  are opposite in the sense that

$$(-\varphi)^{(-f)} = (-f) \Delta (-(-\varphi)) = (-f) \Delta \varphi = f^\varphi. \tag{4}$$

The definition of  $f^\varphi$  is closely connected to the deconvolution operation. For any  $g, h : X \rightarrow \overline{\mathbb{R}}$ , the *deconvolution* of  $g$  and  $h$  is the function  $g \ominus h$  defined by

$$(g \ominus h)(x) = \sup_{y-z=x} (g(y) - h(z)),$$

for every  $x \in X$ . Denoting by  $h_-$  the function defined by  $h_-(x) = h(-x)$  for every  $x \in X$ , we deduce immediately from the above definition that

$$g \ominus h = g \Delta (-h_-) = (h_-)^g. \tag{5}$$

It ensues that for any  $f, \varphi : X \rightarrow \overline{\mathbb{R}}$ ,

$$f^\varphi = \varphi \ominus f_-.$$

The deconvolution operation has been studied in detail by many authors, see, for example, [3, 12, 14, 38].

Following the terminology of Moreau [21], we call  $\varphi$ -*elementary function* a function of the form  $\varphi(\cdot - y) + \lambda$  with  $y \in X$  and  $\lambda \in \mathbb{R}$ . By using a generalized conjugacy argument, one can show that for any  $\varphi, f : X \rightarrow \overline{\mathbb{R}}$ ,

$$(f^{\varphi-})^\varphi \text{ is the upper envelope of the } \varphi\text{-elementary functions that minorize } f, \tag{6}$$

see, for example, [21, §4] and [30, §11.L]. One can easily deduce the following characterization of  $\varphi$ -envelopes: for any  $g : X \rightarrow \overline{\mathbb{R}}$ ,

$$g \text{ is the upper envelope of a family of } \varphi\text{-elementary functions} \tag{7}$$

$$\begin{aligned} &\Updownarrow \\ g &= (g^{\varphi-})^\varphi \tag{8} \end{aligned}$$

$$\begin{aligned} &\Updownarrow \\ g &\text{ is a } \varphi\text{-envelope.} \tag{9} \end{aligned}$$

The expression of the double envelope  $(g^{\varphi-})^\varphi$  can be developed as follows

$$\begin{aligned} (g^{\varphi-})^\varphi &= \varphi \Delta (-g^{\varphi-}) \\ &= \varphi \Delta (-(\varphi_- \Delta (-g))) \\ &= \varphi \Delta ((-\varphi_-) \nabla g). \end{aligned}$$

By using the deconvolution operation, we obtain

$$\begin{aligned} (g^{\varphi-})^\varphi &= \varphi \ominus (\varphi \Delta (-g_-)) \\ &= \varphi \ominus (\varphi \ominus g). \end{aligned}$$

From the equivalence (8)  $\Leftrightarrow$  (9), we deduce that

$$\begin{aligned} g \text{ is a } \varphi\text{-envelope} & \\ \Updownarrow & \\ g = \varphi \Delta ((-\varphi_-) \nabla g) & \tag{10} \\ \Updownarrow & \\ g = \varphi \ominus (\varphi \ominus g). & \end{aligned}$$

Now let  $f, \psi : X \rightarrow \overline{\mathbb{R}}$ . Following the terminology of Martinez-Legaz and Penot [18], the function  $f$  is said to be (exactly)  $\psi$ -regular if  $f = (f \ominus \psi) \nabla \psi$ . By taking the opposite in each member of the equality (10), we find

$$\begin{aligned} -g &= (-\varphi) \nabla (\varphi_- \Delta (-g)) \\ &= (-\varphi) \nabla ((-g) \ominus (-\varphi)). \end{aligned}$$

In view of the above equivalences, this implies that

$$g \text{ is a } \varphi\text{-envelope} \iff -g \text{ is } (-\varphi)\text{-regular in the sense of [18].}$$

We denote by  $\mathcal{E}^\varphi(X)$ , or  $\mathcal{E}^\varphi$  if there is no risk of confusion, the set of  $\varphi$ -envelopes and by  $F_\varphi : \mathcal{E}^{\varphi-} \rightarrow \mathcal{E}^\varphi$  the map defined by  $F_\varphi(f) = f^\varphi$  for every  $f \in \mathcal{E}^{\varphi-}$ . The equivalence (8)  $\Leftrightarrow$  (9) says that  $F_\varphi \circ F_{\varphi-} = \text{Id}_{\mathcal{E}^\varphi}$  and  $F_{\varphi-} \circ F_\varphi = \text{Id}_{\mathcal{E}^{\varphi-}}$ , otherwise stated we have the following proposition.

PROPOSITION 3.1. *The map  $F_\varphi : \mathcal{E}^{\varphi-} \rightarrow \mathcal{E}^\varphi$  is bijective and  $(F_\varphi)^{-1} = F_{\varphi-}$ .*

As a consequence of the previous proposition, if  $\varphi$  is even the map  $F_\varphi : \mathcal{E}^{\varphi-} \rightarrow \mathcal{E}^\varphi$  is bijective and  $(F_\varphi)^{-1} = F_\varphi$ .

Let us now state several general properties of  $\varphi$ -envelopes.

PROPOSITION 3.2. *Let  $X$  be a vector space and let  $\varphi : X \rightarrow \overline{\mathbb{R}}$ .*

- (i) *For every function  $f : X \rightarrow \overline{\mathbb{R}}$  and every  $a \in X$  and  $\beta \in \mathbb{R}$ , we have  $(f(\cdot - a) - \beta)^\varphi = f^\varphi(\cdot - a) + \beta$ . If  $g \in \mathcal{E}^\varphi$ , then  $g(\cdot - a) + \beta \in \mathcal{E}^\varphi$  for every  $a \in X$  and  $\beta \in \mathbb{R}$ .*

- (ii) Given a family  $(f_i)_{i \in I}$  of functions  $f_i : X \rightarrow \overline{\mathbb{R}}$ , we have  $(\inf_{i \in I} f_i)^\varphi = \sup_{i \in I} f_i^\varphi$ . If  $g = \sup_{i \in I} g_i$  with  $g_i \in \mathcal{E}^\varphi$  for every  $i \in I$ , then  $g \in \mathcal{E}^\varphi$ .
- (iii) For  $f_1, f_2 : X \rightarrow \overline{\mathbb{R}}$ , we have  $(f_1 \nabla f_2)^\varphi = f_1^{(f_2^\varphi)}$ . Let  $g, h : X \rightarrow \overline{\mathbb{R}}$ . If  $h \in \mathcal{E}^g$  and  $g \in \mathcal{E}^\varphi$ , then  $h \in \mathcal{E}^\varphi$ . Otherwise stated, if  $g \in \mathcal{E}^\varphi$ , then  $\mathcal{E}^g \subset \mathcal{E}^\varphi$ .
- (iv) For  $f : X \rightarrow \overline{\mathbb{R}}$ , we have  $(f^\varphi)_- = f_-^{\varphi-}$ . As a consequence,  $g \in \mathcal{E}^\varphi$  if and only if  $g_- \in \mathcal{E}^{\varphi-}$ .

*Proof.* (i) Let  $a \in X$  and  $\beta \in \mathbb{R}$ . For every  $x \in X$ , we have

$$\begin{aligned} (f(\cdot - a) - \beta)^\varphi(x) &= \sup_{y \in X} \{\varphi(x - y) \dot{-} f(y - a) + \beta\} \\ &= \sup_{y' \in X} \{\varphi(x - a - y') \dot{-} f(y') + \beta\} = f^\varphi(x - a) + \beta. \end{aligned}$$

For the second assertion of (i), it suffices to apply the first part with  $g = f^\varphi$ .

(ii) By definition, we have

$$\begin{aligned} \left(\inf_{i \in I} f_i\right)^\varphi &= \varphi \Delta \left(-\inf_{i \in I} f_i\right) \\ &= \varphi \Delta \sup_{i \in I} (-f_i) \\ &= \sup_{i \in I} (\varphi \Delta (-f_i)) = \sup_{i \in I} f_i^\varphi, \quad \text{see, for example, [21].} \end{aligned}$$

Now assume that  $g = \sup_{i \in I} g_i$  with  $g_i \in \mathcal{E}^\varphi$  for every  $i \in I$ . Then, for each  $i \in I$ , we have  $g_i = f_i^\varphi$  for some  $f_i$ . It ensues that  $g = \sup_{i \in I} f_i^\varphi = (\inf_{i \in I} f_i)^\varphi$ , hence  $g \in \mathcal{E}^\varphi$ .

(iii) By definition, we have

$$\begin{aligned} f_1^{(f_2^\varphi)} &= f_2^\varphi \Delta (-f_1) \\ &= (\varphi \Delta (-f_2)) \Delta (-f_1) \\ &= \varphi \Delta ((-f_2) \Delta (-f_1)) \\ &= \varphi \Delta (-(f_2 \nabla f_1)) \\ &= (f_2 \nabla f_1)^\varphi = (f_1 \nabla f_2)^\varphi. \end{aligned}$$

Now assume that  $h \in \mathcal{E}^g$  and  $g \in \mathcal{E}^\varphi$ . Then there exist  $f_1, f_2 : X \rightarrow \overline{\mathbb{R}}$  such that  $h = f_1^g$  and  $g = f_2^\varphi$ . It ensues that  $h = f_1^{(f_2^\varphi)} = (f_1 \nabla f_2)^\varphi$ , hence  $h \in \mathcal{E}^\varphi$ .

(iv) For every  $x \in X$ , we have

$$\begin{aligned} (f^\varphi)_-(x) &= \sup_{y \in X} \{\varphi(-x - y) \dot{-} f(y)\} \\ &= \sup_{\xi \in X} \{\varphi(-x + \xi) \dot{-} f(-\xi)\} \\ &= \sup_{\xi \in X} \{\varphi_-(x - \xi) \dot{-} f_-(\xi)\} = f_-^{\varphi-}(x). \end{aligned}$$

If  $g \in \mathcal{E}^\varphi$ , there exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $g = f^\varphi$ . It ensues that  $g_- = (f^\varphi)_- = (f_-)^{\varphi-}$ , hence  $g_- \in \mathcal{E}^{\varphi-}$ . The proof of the reverse assertion is identical.  $\square$



The equalities in assertions (i) and (ii) of the above proposition are also consequences of [32, Theorem 3.1]† giving a general description of generalized conjugacy from  $\mathcal{F}(X, \overline{\mathbb{R}})$  (the set of functions from  $X$  to  $\overline{\mathbb{R}}$ ) into itself. The above proofs are provided for completeness.

In the next proposition, we show that the  $\varphi$ -envelope of a continuous linear functional is affine and we characterize the elements of  $\mathcal{E}^\varphi$  that are linear.

**PROPOSITION 3.3.** *Let  $X$  be a locally convex space. Let  $\varphi : X \rightarrow \overline{\mathbb{R}}$  and  $\xi^* \in X^*$ . Then we have:*

- (i)  $\langle \xi^*, \cdot \rangle^\varphi = -\langle \xi^*, \cdot \rangle + (-\varphi)^*(\xi^*)$ ;
- (ii) if  $\varphi \neq -\omega_X$ , the following equivalence holds

$$\langle \xi^*, \cdot \rangle \in \mathcal{E}^\varphi \iff \xi^* \in -\text{dom}(-\varphi)^*.$$

*Proof.* (i) For every  $x \in X$ , we have

$$\begin{aligned} \langle \xi^*, \cdot \rangle^\varphi(x) &= \sup_{y \in X} \{\varphi(y) - \langle \xi^*, x - y \rangle\} \\ &= -\langle \xi^*, x \rangle + (-\varphi)^*(\xi^*). \end{aligned}$$

(ii) Let  $g = \langle \xi^*, \cdot \rangle$ . We deduce from (i) that

$$g^{\varphi^-} = -\langle \xi^*, \cdot \rangle + (-\varphi_-)^*(\xi^*) = -\langle \xi^*, \cdot \rangle + (-\varphi)^*(-\xi^*). \tag{11}$$

First assume that  $(-\varphi)^*(-\xi^*) = +\infty$ . Then we have  $g^{\varphi^-} = \omega_X$ , thus implying that  $(g^{\varphi^-})^\varphi = -\omega_X$ . It ensues that  $(g^{\varphi^-})^\varphi \neq g$ , which shows that  $g \notin \mathcal{E}^\varphi$  according to the equivalence (7)  $\iff$  (8). Now assume that  $(-\varphi)^*(-\xi^*) < +\infty$ . Observe that  $(-\varphi)^*(-\xi^*) \in \mathbb{R}$  since

$$(-\varphi)^*(-\xi^*) = -\infty \implies \sup_{x \in X} \langle -\xi^*, x \rangle + \varphi(x) = -\infty \implies \varphi = -\omega_X,$$

which is impossible by assumption. Since  $(-\varphi)^*(-\xi^*) \in \mathbb{R}$ , we deduce from (11), (i) above and Proposition 3.2(i) that

$$(g^{\varphi^-})^\varphi = \langle \xi^*, \cdot \rangle + (-\varphi)^*(-\xi^*) - (-\varphi)^*(-\xi^*) = \langle \xi^*, \cdot \rangle = g,$$

and therefore  $g \in \mathcal{E}^\varphi$ . □

For every set  $C \subset X$ , let us set

$$\Sigma_C = \{f : X \rightarrow \overline{\mathbb{R}}, \text{dom } f \subset C\} \quad \text{and} \quad \Sigma_C^* = \{f^*, f \in \Sigma_C\}.$$

We adopt the same notations  $\Sigma_D$  and  $\Sigma_D^*$  for a subset  $D \subset X^*$ .

**THEOREM 3.1.** *Let  $X$  be a locally convex space and let  $\varphi : X \rightarrow \overline{\mathbb{R}}$  be such that  $\varphi \neq -\omega_X$ . For every subset  $D$  of  $X^*$ , the following assertions are equivalent:*

† We thank the referee for pointing out the reference [32] to us.

- (i)  $\Sigma_D^* \subset \mathcal{E}^\varphi$ ;
- (ii)  $\{f \in \Gamma_0(X), \text{dom } f^* \subset D\} \subset \mathcal{E}^\varphi$ ;
- (iii)  $D \subset -\text{dom}(-\varphi)^*$ .

*Proof.* (i)  $\Rightarrow$  (ii) Let  $D \subset X^*$ . Observe that

$$\begin{aligned} \{f \in \Gamma_0(X), \text{dom } f^* \subset D\} &= \{g^*, \text{dom } g \subset D \text{ and } g \in \Gamma_0(X)\} \\ &\subset \{g^*, \text{dom } g \subset D\} = \Sigma_D^*. \end{aligned}$$

The implication (i)  $\Rightarrow$  (ii) follows immediately.

(ii)  $\Rightarrow$  (iii) Assume that

$$\{f \in \Gamma_0(X), \text{dom } f^* \subset D\} \subset \mathcal{E}^\varphi. \tag{12}$$

Let  $\xi^* \in D$ . Observe that  $\langle \xi^*, \cdot \rangle \in \Gamma_0(X)$  and that

$$\text{dom}(\langle \xi^*, \cdot \rangle)^* = \text{dom } \delta_{\{\xi^*\}} = \{\xi^*\} \subset D,$$

hence  $\langle \xi^*, \cdot \rangle \in \mathcal{E}^\varphi$  in view of (12). We then deduce from Proposition 3.3(ii) that  $\xi^* \in -\text{dom}(-\varphi)^*$ . Since this is true for every  $\xi^* \in D$ , we conclude that  $D \subset -\text{dom}(-\varphi)^*$ .

(iii)  $\Rightarrow$  (i) Now assume that  $D \subset -\text{dom}(-\varphi)^*$  and let  $f \in \Sigma_D^*$ . There exists  $g : X^* \rightarrow \overline{\mathbb{R}}$  such that  $f = g^*$  and  $\text{dom } g \subset D$ . The definition of the Legendre–Fenchel conjugate yields

$$\begin{aligned} f &= \sup_{\xi^* \in X^*} \{\langle \xi^*, \cdot \rangle - g(\xi^*)\} \\ &= \sup_{\xi^* \in \text{dom } g} \{\langle \xi^*, \cdot \rangle - g(\xi^*)\}. \end{aligned} \tag{13}$$

Recalling that  $\text{dom } g \subset D \subset -\text{dom}(-\varphi)^*$ , we deduce from Proposition 3.3(ii) that the linear function  $\langle \xi^*, \cdot \rangle$  is a  $\varphi$ -envelope for every  $\xi^* \in \text{dom } g$ . In view of Proposition 3.2(i), the affine function  $\langle \xi^*, \cdot \rangle - g(\xi^*)$  is also a  $\varphi$ -envelope for every  $\xi^* \in \text{dom } g$ . Coming back to formula (13), we infer from Proposition 3.2(ii) that  $f$  is a  $\varphi$ -envelope as a supremum of  $\varphi$ -envelopes. Finally, we have shown that  $f \in \mathcal{E}^\varphi$ , which proves the inclusion  $\Sigma_D^* \subset \mathcal{E}^\varphi$ .  $\square$

Given a set  $D \subset X^*$ , the following result explores the links between the class  $\Sigma_D^*$  and the class of functions  $f \in \Gamma_0(X)$  satisfying  $\text{dom } f^* \subset D$ . When the set  $D$  is  $w(X^*, X)$ -closed and convex, these classes can be characterized via the support function of  $D$ .

**PROPOSITION 3.4.** *Let  $X$  be a locally convex space and let  $D$  be a non-empty subset of  $X^*$ .*

(i) *We have*

$$\{f \in \Gamma_0(X), \text{dom } f^* \subset D\} \cup \{\omega_X, -\omega_X\} \subset \Sigma_D^*, \tag{14}$$

$$\Sigma_D^* \subset \{f \in \Gamma_0(X), \text{dom } f^* \subset \overline{\text{co}}^{w^*}(D)\} \cup \{\omega_X, -\omega_X\}. \tag{15}$$

As a consequence, if the set  $D \subset X^*$  is  $w(X^*, X)$ -closed and convex, the following equality holds true

$$\Sigma_D^* = \{f \in \Gamma_0(X), \text{dom } f^* \subset D\} \cup \{\omega_X, -\omega_X\}. \tag{16}$$

(ii) If the set  $D \subset X^*$  is  $w(X^*, X)$ -closed and convex, then

$$\begin{aligned} & \{f \in \Gamma_0(X), \text{dom } f^* \subset D\} \\ &= \{f \in \Gamma_0(X), f^\infty \leq \sigma_D\} \end{aligned} \tag{17}$$

$$= \{f \in \Gamma_0(X), f(y) \leq f(x) + \sigma_D(y - x), \forall x, y \in X\}. \tag{18}$$

*Proof.* (i) We have already shown the inclusion  $\{f \in \Gamma_0(X), \text{dom } f^* \subset D\} \subset \Sigma_D^*$ , see the proof of Theorem 3.1. On the other hand, we always have  $-\omega_X \in \Sigma_D^*$ . Since  $D \neq \emptyset$ , we also have  $\omega_X \in \Sigma_D^*$ . This proves the inclusion (14). Let us now establish (15). Assume that  $f \in \Sigma_D^*$ . There exists  $g : X^* \rightarrow \overline{\mathbb{R}}$  such that  $\text{dom } g \subset D$  and  $f = g^*$ . We distinguish the cases  $\overline{\text{co}}^{w^*}(g)$  proper and  $\overline{\text{co}}^{w^*}(g)$  improper. If  $\overline{\text{co}}^{w^*}(g) = \omega_{X^*}$ , we have  $g = \omega_{X^*}$ , hence  $f = -\omega_X$ . If  $\overline{\text{co}}^{w^*}(g)$  takes the value  $-\infty$ , we infer that  $g^* = (\overline{\text{co}}^{w^*}(g))^* = \omega_X$ , whence  $f = \omega_X$ . Let us now assume that  $\overline{\text{co}}^{w^*}(g) \in \Gamma_0(X^*)$ . It ensues that  $f = g^* = (\overline{\text{co}}^{w^*}(g))^* \in \Gamma_0(X)$ . This implies in turn that  $f^* = \overline{\text{co}}^{w^*}(g)$ , thus

$$\text{dom } f^* = \text{dom}(\overline{\text{co}}^{w^*}(g)) \subset \overline{\text{co}}^{w^*}(\text{dom } g) \subset \overline{\text{co}}^{w^*}(D),$$

which ends the proof of (15). When the set  $D$  is  $w(X^*, X)$ -closed and convex, equality (16) is an immediate consequence of the inclusions (14)–(15).

(ii) Assuming that the set  $D$  is  $w(X^*, X)$ -closed and convex, we have  $\text{dom } f^* \subset D$  if and only if  $\sigma_{\text{dom } f^*} \leq \sigma_D$ . Recalling that  $\sigma_{\text{dom } f^*} = f^\infty$  (see §2), we derive equality (17). Since  $f^\infty = \sup_{x \in \text{dom } f} (f(\cdot + x) - f(x))$ , we deduce in turn equality (18).  $\square$

*Remark 3.1.* In general, the inclusions (14) and (15) are strict, as will be shown in Example 7.1.

If  $X$  is a Banach space and if the set  $D \subset X^*$  is closed, the class of functions  $f \in \Gamma_0(X)$  satisfying  $\text{dom } f^* \subset D$  can be expressed via the subdifferential of  $f$ .

**PROPOSITION 3.5.** *Let  $X$  be a Banach space and let  $D$  be a closed subset of  $X^*$ . Then we have*

$$\{f \in \Gamma_0(X), \text{dom } f^* \subset D\} = \{f \in \Gamma_0(X), \partial f(x) \subset D \text{ for all } x \in X\}.$$

*Proof.* Let us first state as a lemma the following direct consequence of the Brønsted–Rockafellar theorem (see [5, Theorem 2]) concerning the conjugate of a function in  $\Gamma_0(X)$ .

**LEMMA 3.1** (See [5, Theorem 2]). *If  $X$  is a Banach space and if  $f \in \Gamma_0(X)$ , then  $\text{cl}(\text{dom } f^*) = \text{cl}(\text{Rge}(\partial f))$ .*

Assume that the set  $D \subset X^*$  is closed. From Lemma 3.1, we have for every  $f \in \Gamma_0(X)$ ,

$$\begin{aligned} \text{dom } f^* \subset D &\iff \text{Rge}(\partial f) \subset D \\ &\iff \partial f(x) \subset D \quad \text{for all } x \in X. \end{aligned}$$

The announced equality follows immediately. □

Applying Theorem 3.1 with particular sets  $D$ , we obtain the following corollaries.

**COROLLARY 3.1.** *Let  $X$  be a locally convex space and let  $\varphi : X \rightarrow \overline{\mathbb{R}}$  be such that  $\varphi \neq -\omega_X$ . Then the following equivalence holds:*

$$\Gamma(X) \subset \mathcal{E}^\varphi \iff \text{dom}(-\varphi)^* = X^*.$$

*Proof.* It suffices to take  $D = X^*$  in the equivalence (i)  $\Leftrightarrow$  (iii) of Theorem 3.1. □

*Remark 3.2.* Under the assumption  $\text{dom}(-\varphi)^* = X^*$ , the function  $\varphi$  cannot be convex (see hereafter). Therefore, the set  $\mathcal{E}^\varphi$  is strictly larger than  $\Gamma(X)$ , since it contains the non-convex function  $\varphi$ .

If  $\text{dom}(-\varphi)^* = X^*$ , we have  $(-\varphi)^*(0) < +\infty$ . Recalling that  $(-\varphi)^*(0) = \sup \varphi$ , we deduce that the function  $\varphi$  is bounded from above on the whole space  $X$ . If, moreover, the function  $\varphi$  is convex, we infer from a classical result that it is constant, say  $\varphi \equiv \beta$  for some  $\beta \in \mathbb{R}$ . It ensues that  $(-\varphi)^* = \beta + \delta_{\{0\}}$ , hence  $\text{dom}(-\varphi)^* = \{0\}$ , a contradiction. This confirms that functions  $\varphi$  with  $\text{dom}(-\varphi)^* = X^*$  cannot be convex.

Given a set  $K \subset X$ , recall that a function  $f : X \rightarrow \mathbb{R} \cup \{+\infty\}$  is said to be  $K$ -non-increasing (respectively  $K$ -non-decreasing) if  $f(y) \leq f(x)$  (respectively  $f(y) \geq f(x)$ ) for all  $x, y \in X$  such that  $y - x \in K$ .

**COROLLARY 3.2.** *Let  $X$  be a locally convex space. Let  $K \subset X$  be a closed convex cone and let  $\varphi : X \rightarrow \overline{\mathbb{R}}$  be such that  $\varphi \neq -\omega_X$ . Then the set  $\mathcal{E}^\varphi$  contains all the functions of  $\Gamma_0(X)$  which are  $K$ -non-increasing if and only if  $-K^\circ \subset \text{dom}(-\varphi)^*$ .*

*Proof.* Take  $D = K^\circ$  in the equivalence (ii)  $\Leftrightarrow$  (iii) of Theorem 3.1 to obtain that

$$\begin{aligned} \{f \in \Gamma_0(X), \text{dom } f^* \subset K^\circ\} \subset \mathcal{E}^\varphi &\iff K^\circ \subset -\text{dom}(-\varphi)^* \\ &\iff -K^\circ \subset \text{dom}(-\varphi)^*. \end{aligned} \tag{19}$$

On the other hand, observe by (18) that for  $f \in \Gamma_0(X)$ ,

$$\begin{aligned} \text{dom } f^* \subset K^\circ &\iff f(y) \leq f(x) + \sigma_{K^\circ}(y - x) \quad \text{for all } x, y \in X \\ &\iff f(y) \leq f(x) + \delta_K(y - x) \quad \text{for all } x, y \in X \\ &\iff f \text{ is } K\text{-non-increasing.} \end{aligned} \tag{20}$$

The announced equivalence then follows immediately from (19) and (20). □

In the following, when  $X$  is a normed space we will denote by  $\mathbb{B}_X$  (respectively  $\mathbb{B}_{X^*}$ ) the closed unit ball of  $X$  (respectively  $X^*$ ).

**COROLLARY 3.3.** *Let  $(X, \|\cdot\|)$  be a normed space. Let a real  $k \geq 0$  and let  $\varphi : X \rightarrow \overline{\mathbb{R}}$  be such that  $\varphi \neq -\omega_X$ . Then the set  $\mathcal{E}^\varphi$  contains all the functions of  $\Gamma_0(X)$  which are  $k$ -Lipschitz continuous on  $X$  if and only if  $k\mathbb{B}_{X^*} \subset \text{dom}(-\varphi)^*$ .*

*Proof.* Take  $D = k\mathbb{B}_{X^*}$  in the equivalence (ii)  $\Leftrightarrow$  (iii) of Theorem 3.1 to obtain that

$$\begin{aligned} \{f \in \Gamma_0(X), \text{dom } f^* \subset k\mathbb{B}_{X^*}\} \subset \mathcal{E}^\varphi &\iff k\mathbb{B}_{X^*} \subset -\text{dom}(-\varphi)^* \\ &\iff k\mathbb{B}_{X^*} \subset \text{dom}(-\varphi)^*. \end{aligned} \tag{21}$$

Then observe by (18) that for  $f \in \Gamma_0(X)$ ,

$$\begin{aligned} \text{dom } f^* \subset k\mathbb{B}_{X^*} &\iff f(y) \leq f(x) + k\|y - x\| \quad \text{for all } x, y \in X \\ &\iff f \text{ is } k\text{-Lipschitz on } X, \end{aligned} \tag{22}$$

where the last equivalence is obtained by reversing the roles of  $x$  and  $y$ . The announced equivalence then follows immediately from (21) and (22).  $\square$

§4. *Equivalence between functions and sets.* Recall that for  $f : X \rightarrow \overline{\mathbb{R}}$ , the epigraph (respectively hypograph) of  $f$  is defined by

$$\begin{aligned} \text{epi } f &= \{(x, \lambda) \in X \times \mathbb{R}, f(x) \leq \lambda\} \\ \text{(respectively hypo } f &= \{(x, \lambda) \in X \times \mathbb{R}, f(x) \geq \lambda\}). \end{aligned}$$

The following result allows us to express the epigraph of  $f^\varphi$  as an intersection of sets which are translated from the epigraph of  $\varphi$ .

**PROPOSITION 4.1.** *Let  $X$  be a vector space and let  $\varphi : X \rightarrow \overline{\mathbb{R}}$ .*

(i) *For every  $f : X \rightarrow \overline{\mathbb{R}}$ , we have*

$$\text{epi } f^\varphi = \bigcap_{u \in \text{hypo}(-f)} u + \text{epi } \varphi.$$

(ii) *For every  $g : X \rightarrow \overline{\mathbb{R}}$ , the following equivalence holds:*

$$g \in \mathcal{E}^\varphi \iff \text{epi } g = \bigcap_{u \in U} u + \text{epi } \varphi \quad \text{for some } U \subset X \times \mathbb{R}.$$

*Proof.* (i) For every  $x \in X$  and  $\lambda \in \mathbb{R}$ , the following equivalences hold true:

$$\begin{aligned} (x, \lambda) \in \text{epi } f^\varphi &\iff f^\varphi(x) \leq \lambda \\ &\iff \varphi(x - y) + f(y) \leq \lambda \quad \text{for every } y \in X \\ &\iff \varphi(x - y) - \mu \leq \lambda \quad \text{for every } (y, \mu) \in \text{epi } f \\ &\iff (x - y, \lambda + \mu) \in \text{epi } \varphi \quad \text{for every } (y, \mu) \in \text{epi } f \\ &\iff (x, \lambda) \in (y, -\mu) + \text{epi } \varphi \quad \text{for every } (y, \mu) \in \text{epi } f \\ &\iff (x, \lambda) \in u + \text{epi } \varphi \quad \text{for every } u \in \text{hypo}(-f). \end{aligned}$$

The announced equality follows.

(ii) Let  $g : X \rightarrow \overline{\mathbb{R}}$ . If  $g \in \mathcal{E}^\varphi$ , there exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $g = f^\varphi$ . In view of (i), we obtain that  $\text{epi } g = \bigcap_{u \in U} u + \text{epi } \varphi$  with  $U = \text{hypo}(-f)$ . Conversely, assume that  $\text{epi } g = \bigcap_{u \in U} u + \text{epi } \varphi$  for some  $U \subset X \times \mathbb{R}$ . Then we have

$$\begin{aligned} \text{epi } g &= \bigcap_{(x,\lambda) \in U} (x, \lambda) + \text{epi } \varphi \\ &= \bigcap_{(x,\lambda) \in U} \text{epi}[\varphi(\cdot - x) + \lambda] \\ &= \text{epi} \left[ \sup_{(x,\lambda) \in U} \varphi(\cdot - x) + \lambda \right]. \end{aligned}$$

Hence, we deduce that  $g = \sup_{(x,\lambda) \in U} (\varphi(\cdot - x) + \lambda)$ , which shows by (i) and (ii) in Proposition 3.2 that  $g \in \mathcal{E}^\varphi$ . □

Given a set  $\Lambda \subset X$ , the previous result suggests to consider the class  $\mathcal{I}^\Lambda$  of subsets of  $X$  defined as follows†

$$\mathcal{I}^\Lambda = \{C^\Lambda, C \subset X\}, \quad \text{where } C^\Lambda = \bigcap_{x \in C} x + \Lambda.$$

By convention‡, we take  $\emptyset^\Lambda = \bigcap_{x \in \emptyset} x + \Lambda = X$  for every set  $\Lambda \subset X$ . This implies that  $X \in \mathcal{I}^\Lambda$  for every  $\Lambda \subset X$ . It is immediate to check that  $\mathcal{I}^X = \{X\}$ , while  $\mathcal{I}^\emptyset = \{\emptyset, X\}$ . A set  $D \subset X$  belongs to the class  $\mathcal{I}^\Lambda$  if it is equal to some intersection of translated sets from  $\Lambda$ . It ensues immediately that the class  $\mathcal{I}^\Lambda$  is stable under translation and intersection.

*Example 4.1.* Take  $r > 0$  and  $\Lambda = r\mathbb{B}_X$ . The class  $\mathcal{I}^{r\mathbb{B}_X}$  corresponds to the class studied by Vial [36] under the terminology of  $r$ -strongly convex sets. More generally, for a closed convex set  $\Lambda \subset X$ , the sets of the form  $C^\Lambda$  are called  $\Lambda$ -strongly convex. The  $\Lambda$ -strongly convex sets are thoroughly studied by Polovinkin [25], under an additional condition on the set  $\Lambda$  (which is assumed to be generating, see [25] for more details).

The definition of  $C^\Lambda$  is directly linked to the star-difference of sets. For every  $C_1, C_2 \subset X$ , the star-difference of  $C_1$  with  $C_2$  is the set  $C_1 \overset{*}{-} C_2$  given by

$$C_1 \overset{*}{-} C_2 = \bigcap_{x \in C_2} C_1 - x.$$

We deduce immediately from the above definition that  $C^\Lambda = \Lambda \overset{*}{-} (-C)$  for every  $C, \Lambda \subset X$ . The star-difference of sets was used in [27] in the context of differential games. See also [12] for the links between the star-difference of sets and the deconvolution operation, also called epigraphical star-difference.

Given  $C \subset X$  and  $\Lambda \subset X$ , the next proposition gives several expressions for the set  $C^\Lambda$ .

† We draw the attention of the reader to the fact that the notation  $C^\Lambda$  must not be confused with that of the set of maps from  $\Lambda$  into  $C$ .

‡ In particular, we obtain  $\emptyset^\emptyset = X$ .

PROPOSITION 4.2. *Let  $X$  be a vector space. For any sets  $C \subset X$  and  $\Lambda \subset X$ , we have:*

- (i)  $C^\Lambda = \{x \in X, x - C \subset \Lambda\} = \{x \in X, C \subset x - \Lambda\}$ ;
- (ii)  $X \setminus C^\Lambda = C + (X \setminus \Lambda)$  or equivalently  $C^{X \setminus \Lambda} = X \setminus (C + \Lambda)$ ;
- (iii)  $(X \setminus \Lambda)^{X \setminus C} = C^\Lambda$ .

*Proof.* (i) It suffices to observe that

$$\begin{aligned} x \in C^\Lambda &\iff \text{for all } u \in C, x \in u + \Lambda \\ &\iff \text{for all } u \in C, x - u \in \Lambda \\ &\iff x - C \subset \Lambda \\ &\iff C \subset x - \Lambda. \end{aligned}$$

(ii) From the definition of  $C^\Lambda$ , we deduce immediately that

$$X \setminus C^\Lambda = \bigcup_{u \in C} u + (X \setminus \Lambda) = C + (X \setminus \Lambda),$$

which is the first equality in (ii). From this equality with  $X \setminus \Lambda$  in place of  $\Lambda$ , we obtain that  $X \setminus C^{X \setminus \Lambda} = C + \Lambda$ , or equivalently  $C^{X \setminus \Lambda} = X \setminus (C + \Lambda)$ .

(iii) We infer from the previous assertion that

$$X \setminus [(X \setminus \Lambda)^{X \setminus C}] = (X \setminus \Lambda) + C = X \setminus C^\Lambda,$$

whence the equality  $(X \setminus \Lambda)^{X \setminus C} = C^\Lambda$ . □

The elements  $D$  of  $\mathcal{I}^\Lambda$  can be characterized by the equality  $(D^{-\Lambda})^\Lambda = D$ . This is the subject of the next proposition.

PROPOSITION 4.3. *Let  $X$  be a vector space and let  $\Lambda \subset X$ . For any set  $D \subset X$ , the set  $(D^{-\Lambda})^\Lambda$  is the smallest element of  $\mathcal{I}^\Lambda$  containing the set  $D$ . As a consequence, the following equivalence holds true:*

$$D \in \mathcal{I}^\Lambda \iff (D^{-\Lambda})^\Lambda = D.$$

*Proof.* Let  $S$  be the subset of  $X$  defined by

$$S = \bigcap_{x \in X, x + \Lambda \supset D} x + \Lambda.$$

We clearly have  $S \in \mathcal{I}^\Lambda$  and  $S \supset D$ . Now let any  $S' \in \mathcal{I}^\Lambda$  with  $S' \supset D$ . By definition, there exists some  $C \subset X$  such that  $S' = \bigcap_{x \in C} x + \Lambda$ . The inclusion  $S' \supset D$  implies that  $x + \Lambda \supset D$  for every  $x \in C$  and therefore

$$S' = \bigcap_{x \in C} x + \Lambda \supset \bigcap_{x \in X, x + \Lambda \supset D} x + \Lambda = S.$$

This proves that the set  $S$  is the smallest element of  $\mathcal{I}^\Lambda$  containing  $D$ . Recall now from Proposition 4.2(i) that condition  $x + \Lambda \supset D$  is equivalent to  $x \in D^{-\Lambda}$ . We deduce that

$$S = \bigcap_{x \in D^{-\Lambda}} x + \Lambda = (D^{-\Lambda})^\Lambda.$$

This finishes the proof of the first assertion. The second assertion is an immediate consequence of the first. □

Let us write the expression of the double envelope  $(D^{-\Lambda})^\Lambda$  by using the star-difference operation

$$\begin{aligned} (D^{-\Lambda})^\Lambda &= \Lambda \overset{*}{-} (-(D^{-\Lambda})) \\ &= \Lambda \overset{*}{-} ((-D)^\Lambda) \\ &= \Lambda \overset{*}{-} (\Lambda \overset{*}{-} D). \end{aligned} \tag{23}$$

In view of Proposition 4.2, the complement of the set  $(D^{-\Lambda})^\Lambda$  can be expressed as

$$\begin{aligned} X \setminus (D^{-\Lambda})^\Lambda &= D^{-\Lambda} + X \setminus \Lambda \\ &= -(X \setminus \Lambda)^{X \setminus D} + X \setminus \Lambda \\ &= ((X \setminus D) \overset{*}{-} (X \setminus \Lambda)) + X \setminus \Lambda. \end{aligned} \tag{24}$$

From equalities (23)–(24) and Proposition 4.3, we deduce that

$$\begin{aligned} D &\in \mathcal{I}^\Lambda \\ &\Updownarrow \\ D &= \Lambda \overset{*}{-} (\Lambda \overset{*}{-} D) \\ &\Updownarrow \\ X \setminus D &= ((X \setminus D) \overset{*}{-} (X \setminus \Lambda)) + X \setminus \Lambda. \end{aligned}$$

The last equality amounts to saying that the set  $X \setminus D$  is exactly  $(X \setminus \Lambda)$ -regular in the sense of [18].

With the notation introduced above, for  $f, g : X \rightarrow \overline{\mathbb{R}}$ , the results of Proposition 4.1 can be restated as

$$\text{epi } f^\varphi = (\text{hypo}(-f))^{\text{epi } \varphi}$$

and

$$g \in \mathcal{E}^\varphi \iff \text{epi } g \in \mathcal{I}^{\text{epi } \varphi}.$$

This shows that the study of  $\varphi$ -envelopes amounts to that of the class  $\mathcal{I}^{\text{epi } \varphi}$ . Conversely, given a set  $\Lambda \subset X$ , the class  $\mathcal{I}^\Lambda$  can be fully described via the  $\delta_\Lambda$ -envelopes.



PROPOSITION 4.4. *Let  $X$  be a vector space and let  $\Lambda \subset X$ .*

(i) *For every function  $f : X \rightarrow \overline{\mathbb{R}}$ , we have†*

$$f^{\delta_\Lambda} = -\inf_X f + \delta_{(\text{dom } f)^\Lambda}. \tag{25}$$

*As a consequence, the equality  $(\delta_C)^{\delta_\Lambda} = \delta_{C^\Lambda}$  holds for any non-empty set  $C \subset X$ .*

(ii) *For every function  $g : X \rightarrow \overline{\mathbb{R}}$  such that  $g \neq \pm\omega_X$ , we have*

$$g \in \mathcal{E}^{\delta_\Lambda} \iff g = \beta + \delta_{C^\Lambda} \text{ for some } \beta \in \mathbb{R} \text{ and some } C \neq \emptyset.$$

*Proof.* (i) For every function  $f : X \rightarrow \overline{\mathbb{R}}$  and every  $x \in X$ , the definition of  $f^{\delta_\Lambda}$  gives

$$f^{\delta_\Lambda}(x) = \sup_{y \in X} \{\delta_\Lambda(x - y) \dot{-} f(y)\} = \sup_{y \in \text{dom } f} \{\delta_\Lambda(x - y) - f(y)\}.$$

First assume that  $x - \text{dom } f \subset \Lambda$ . For every  $y \in \text{dom } f$ , we then have  $x - y \in \Lambda$ , whence  $\delta_\Lambda(x - y) = 0$ . It ensues that

$$f^{\delta_\Lambda}(x) = \sup_{y \in \text{dom } f} -f(y) = \sup_X(-f) = -\inf_X f.$$

Now assume that  $x - \text{dom } f \not\subset \Lambda$ . In this case, there exists  $y \in \text{dom } f$  such that  $x - y \notin \Lambda$ . We then have  $\delta_\Lambda(x - y) = +\infty$ , whence  $f^{\delta_\Lambda}(x) = +\infty$ . Finally, we obtain for every  $x \in X$ ,

$$f^{\delta_\Lambda}(x) = \begin{cases} -\inf_X f & \text{if } x - \text{dom } f \subset \Lambda, \\ +\infty & \text{otherwise.} \end{cases}$$

Condition  $x - \text{dom } f \subset \Lambda$  is equivalent to  $x \in (\text{dom } f)^\Lambda$  in view of Proposition 4.2(i). Formula (25) follows immediately. For the last assertion, it suffices to take  $f = \delta_C$ .

(ii) Let  $g \in \mathcal{E}^{\delta_\Lambda}$  be such that  $g \neq \pm\omega_X$ . There exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $g = f^{\delta_\Lambda}$ , hence we deduce from (i) that  $g = -\inf_X f + \delta_{(\text{dom } f)^\Lambda}$ . Since  $g \neq \pm\omega_X$ , we have  $\inf_X f \in \mathbb{R}$  and  $\text{dom } f \neq \emptyset$ . It suffices then to take  $\beta = -\inf_X f$  and  $C = \text{dom } f$ . Conversely, assume that  $g = \beta + \delta_{C^\Lambda}$  for some  $\beta \in \mathbb{R}$  and some  $C \neq \emptyset$ . Assertion (i) then shows that  $g = f^{\delta_\Lambda}$  for the function  $f$  defined by  $f = -\beta + \delta_C$ , hence  $g \in \mathcal{E}^{\delta_\Lambda}$ . □

*Remark 4.1.* The previous proposition shows that for every  $C, \Lambda \subset X$  with  $C \neq \emptyset$ ,

$$(\delta_C)^{\delta_\Lambda} = (-\delta_C) \triangle \delta_\Lambda = \delta_{C^\Lambda}. \tag{26}$$

† If  $\inf_X f = +\infty$  we have  $\text{dom } f = \emptyset$ , hence  $(\text{dom } f)^\Lambda = X$  and  $\delta_{(\text{dom } f)^\Lambda} \equiv 0$ . Therefore, the addition in the right-hand side of (25) is well-defined.

It is interesting to compare this formula with the following one

$$(\delta_C)^{-\delta_\Lambda} = (-\delta_C) \Delta (-\delta_\Lambda) = -\delta_{C+\Lambda}, \tag{27}$$

that is obtained as a consequence of the equality  $\delta_{C+\Lambda} = \delta_C \nabla \delta_\Lambda$ .

**COROLLARY 4.1.** *Let  $X$  be a vector space. For every set  $\Lambda \subset X$  and every set  $D \subset X$  such that  $D \neq \emptyset$  and  $D \neq X$ , the following equivalence holds:*

$$\delta_D \in \mathcal{E}^{\delta_\Lambda} \iff D \in \mathcal{I}^\Lambda.$$

*In fact, the implication from the left to the right is true as soon as  $D \neq \emptyset$ , while the reverse one is true if  $D \neq X$ .*

*Proof.* First assume that  $\delta_D \in \mathcal{E}^{\delta_\Lambda}$  and that  $D \neq \emptyset$ . There exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $\delta_D = f^{\delta_\Lambda}$ , hence we deduce from Proposition 4.4(i) that  $\delta_D = -\inf_X f + \delta_{(\text{dom } f)^\Lambda}$ . Since  $D \neq \emptyset$ , we have  $\inf_X f = 0$  and  $D = (\text{dom } f)^\Lambda \in \mathcal{I}^\Lambda$ . Conversely, assume that  $D \in \mathcal{I}^\Lambda$  and that  $D \neq X$ . This implies that  $D = C^\Lambda$  for some  $C \neq \emptyset$ , and hence by Proposition 4.4(i) again  $\delta_D = \delta_{C^\Lambda} = (\delta_C)^{\delta_\Lambda} \in \mathcal{E}^{\delta_\Lambda}$ . □

Let us now study the class  $\mathcal{E}^{-\delta_\Lambda}$ . From the generalized conjugation point of view, the case  $\varphi = -\delta_\Lambda$  is a special instance of a coupling functional  $c : X \times Y \rightarrow \overline{\mathbb{R}}$  of the type  $c = -\delta_G$ , where  $G$  is a subset of  $X \times Y$ . The corresponding conjugation operator, which arises in quasiconvex analysis, has been considered in many papers, see, for example, [17, 33, 37].

**PROPOSITION 4.5.** *Let  $X$  be a vector space. Let  $\Lambda$  be a non-empty subset of  $X$  and let  $f : X \rightarrow \overline{\mathbb{R}}$ . Then:*

(i) *we have*

$$f \in \mathcal{E}^{-\delta_\Lambda} \iff f = \sup_{y \in \Lambda} \inf_{z \in \Lambda} f(\cdot - y + z); \tag{28}$$

*this means equivalently that for every  $x \in X$  and every  $\lambda < f(x)$ , there exists  $y \in \Lambda$  such that  $f(x - y + z) \geq \lambda$  for every  $z \in \Lambda$ ;*

(ii) *if  $f \in \mathcal{E}^{-\delta_\Lambda}$  and if  $\Lambda + \Lambda \subset \Lambda$ , then  $f$  is  $\Lambda$ -non-decreasing; conversely, if  $f$  is  $\Lambda$ -non-decreasing and if  $0 \in \Lambda$ , then  $f \in \mathcal{E}^{-\delta_\Lambda}$ .*

*Proof.* (i) The equivalence (7)  $\iff$  (8) yields

$$f \in \mathcal{E}^{-\delta_\Lambda} \iff f = (f^{(-\delta_\Lambda)-})^{-\delta_\Lambda}.$$

On the other hand, we have

$$f^{(-\delta_\Lambda)-} = \sup_{\xi \in X} -\delta_\Lambda(-\xi) \dot{-} f(\cdot - \xi) = \sup_{-\xi \in \Lambda} -f(\cdot - \xi) = - \inf_{z \in \Lambda} f(\cdot + z)$$

and, hence,

$$(f^{(-\delta_\Lambda)_-})^{-\delta_\Lambda} = \sup_{y \in \Lambda} -f^{(-\delta_\Lambda)_-}(\cdot - y) = \sup_{y \in \Lambda} \inf_{z \in \Lambda} f(\cdot - y + z).$$

We deduce immediately the equivalence (28).

Since the inequality  $(f^{(-\delta_\Lambda)_-})^{-\delta_\Lambda} \leq f$  is always satisfied, we infer that  $f \in \mathcal{E}^{-\delta_\Lambda}$  if and only if for every  $x \in X$ ,

$$\sup_{y \in \Lambda} \inf_{z \in \Lambda} f(x - y + z) \geq f(x).$$

The last assertion of (i) follows immediately.

(ii) Assume that  $f \in \mathcal{E}^{-\delta_\Lambda}$  and that  $\Lambda + \Lambda \subset \Lambda$ . Let  $\xi \in \Lambda$ . In view of (28), we have for every  $x \in X$ ,

$$\begin{aligned} f(x + \xi) &= \sup_{y \in \Lambda} \inf_{z \in \Lambda} f(x + \xi - y + z) \\ &= \sup_{y \in \Lambda} \inf_{z' \in \xi + \Lambda} f(x - y + z') \\ &\geq \sup_{y \in \Lambda} \inf_{z' \in \Lambda} f(x - y + z') \quad \text{since } \xi + \Lambda \subset \Lambda \\ &= f(x). \end{aligned}$$

Since this is true for every  $\xi \in \Lambda$ , we infer that  $f$  is  $\Lambda$ -non-decreasing.

Conversely, assume that  $f$  is  $\Lambda$ -non-decreasing and that  $0 \in \Lambda$ . For every  $y, z \in \Lambda$ , we have

$$f(\cdot - y) \leq f(\cdot - y + z) \leq f(\cdot + z).$$

It ensues immediately that

$$\sup_{y \in \Lambda} f(\cdot - y) \leq \sup_{y \in \Lambda} \inf_{z \in \Lambda} f(\cdot - y + z) \leq \inf_{z \in \Lambda} f(\cdot + z).$$

Since  $0 \in \Lambda$ , we obtain  $\sup_{y \in \Lambda} f(\cdot - y) = \inf_{z \in \Lambda} f(\cdot + z) = f$ , and hence  $f = \sup_{y \in \Lambda} \inf_{z \in \Lambda} f(\cdot - y + z)$ . In view of (28), we conclude that  $f \in \mathcal{E}^{-\delta_\Lambda}$ . □

*Remark 4.2.* When  $\Lambda + \Lambda \subset \Lambda$  and  $0 \in \Lambda$ , the equivalence

$$f \in \mathcal{E}^{-\delta_\Lambda} \iff f \text{ is } \Lambda\text{-non-decreasing}$$

can be recovered by using the subadditivity of the function  $\delta_\Lambda$ , see §6.

For a function  $f : X \rightarrow \overline{\mathbb{R}}$  and  $r \in \overline{\mathbb{R}}$ , the notation  $[f \geq r]$  (respectively  $[f > r]$ ) denotes the set  $\{x \in X, f(x) \geq r\}$  (respectively  $\{x \in X, f(x) > r\}$ ). We adopt the corresponding notation for the sublevel sets.

PROPOSITION 4.6. *Given a vector space  $X$  and  $\Lambda \subset X$ , let  $f, g : X \rightarrow \overline{\mathbb{R}}$  be extended real-valued functions.*

(i) *For every  $\lambda \in \overline{\mathbb{R}}$ , we have*

$$[f^{-\delta_\Lambda} \leq \lambda] = [f < -\lambda]^{X \setminus \Lambda}.$$

(ii) *Assume that for every  $\lambda \in \overline{\mathbb{R}}$ , there exists  $C_\lambda \subset X$  such that*

$$[g \leq \lambda] = C_\lambda^{X \setminus \Lambda}.$$

*Then we have  $g = (-h)^{-\delta_\Lambda}$  for the function  $h : X \rightarrow \overline{\mathbb{R}}$  defined by  $h(x) = \sup\{\lambda \in \overline{\mathbb{R}}, x \in C_\lambda\}$ .*

(iii) *The following assertions are equivalent:*

- (a)  $g \in \mathcal{E}^{-\delta_\Lambda}$ ;
- (b) for all  $\lambda \in \overline{\mathbb{R}}$ ,  $[g \leq \lambda] \in \mathcal{I}^{X \setminus \Lambda}$ .

*Proof.* (i) For every  $\lambda \in \overline{\mathbb{R}}$  and  $x \in X$ , the following equivalences hold:

$$\begin{aligned} f^{-\delta_\Lambda}(x) \leq \lambda &\iff \text{for all } y \in X, -\delta_\Lambda(x - y) \dashv f(y) \leq \lambda \\ &\iff \text{for all } y \in X, x - y \in \Lambda \implies -f(y) \leq \lambda \\ &\iff \text{for all } y \in X, -f(y) > \lambda \implies x - y \in X \setminus \Lambda \\ &\iff \text{for all } y \in [f < -\lambda], \quad x \in y + X \setminus \Lambda \\ &\iff x \in [f < -\lambda]^{X \setminus \Lambda}. \end{aligned}$$

We deduce the equality  $[f^{-\delta_\Lambda} \leq \lambda] = [f < -\lambda]^{X \setminus \Lambda}$ .

(ii) Fix  $x \in X$  and observe that

$$g(x) = \sup\{\lambda \in \overline{\mathbb{R}}, g(x) > \lambda\}. \tag{29}$$

By assumption, we have for every  $\lambda \in \overline{\mathbb{R}}$ ,

$$[g \leq \lambda] = \bigcap_{y \in C_\lambda} y + X \setminus \Lambda,$$

hence  $[g > \lambda] = \bigcup_{y \in C_\lambda} y + \Lambda$ . From (29), we infer that

$$\begin{aligned} g(x) &= \sup\{\lambda \in \overline{\mathbb{R}}, \exists y \in C_\lambda, x \in y + \Lambda\} \\ &= \sup\{\lambda \in \overline{\mathbb{R}}, \exists y \in x - \Lambda, y \in C_\lambda\} \\ &= \sup\left(\bigcup_{y \in x - \Lambda} \{\lambda \in \overline{\mathbb{R}}, y \in C_\lambda\}\right) \\ &= \sup_{y \in x - \Lambda} (\sup\{\lambda \in \overline{\mathbb{R}}, y \in C_\lambda\}) \\ &= \sup_{y \in x - \Lambda} h(y) \\ &= \sup_{y \in X} (-\delta_\Lambda(x - y) \dashv h(y)) = (-h)^{-\delta_\Lambda}(x). \end{aligned}$$

Since this is true for every  $x \in X$ , we conclude that  $g = (-h)^{-\delta_\Lambda}$ .

(iii) The implication (a)  $\implies$  (b) follows immediately from (i), while the converse implication is a consequence of (ii).  $\square$

**COROLLARY 4.2.** *Let  $X$  be a vector space and let  $\Lambda \subset X$ . Assume that the function  $g : X \rightarrow \overline{\mathbb{R}}$  is such that  $g(X) = \{\alpha_1, \dots, \alpha_n\} \subset \overline{\mathbb{R}}$ . If  $\inf g = -\infty$ , the assertions (a) and (b) of Proposition 4.6(iii) are equivalent to the following:*

(b') for all  $i \in \{1, \dots, n\}$ ,  $[g \leq \alpha_i] \in \mathcal{I}^{X \setminus \Lambda}$ ;

while if  $\inf g > -\infty$ , they are equivalent to:

(b'') for all  $i \in \{1, \dots, n\}$ ,  $[g \leq \alpha_i] \in \mathcal{I}^{X \setminus \Lambda}$  and  $\Lambda \neq \emptyset$ .

*Proof.* If  $\inf g = -\infty$ , assertion (b) of Proposition 4.6(iii) is clearly equivalent to (b'), while if  $\inf g > -\infty$ , one has to add the complementary condition  $\emptyset \in \mathcal{I}^{X \setminus \Lambda}$ . Then observe that

$$\emptyset \in \mathcal{I}^{X \setminus \Lambda} \iff X^{X \setminus \Lambda} = \emptyset \iff X + \Lambda = X \iff \Lambda \neq \emptyset,$$

and hence the equivalence (b)  $\iff$  (b'') follows.  $\square$

**COROLLARY 4.3.** *Given a vector space  $X$ , let  $\Lambda \subset X$  and  $D \subset X$ . The following equivalences hold:*

- (i)  $-\delta_D \in \mathcal{E}^{-\delta_\Lambda} \iff X \setminus D \in \mathcal{I}^{X \setminus \Lambda}$ ;
- (ii)  $\delta_D \in \mathcal{E}^{-\delta_\Lambda} \iff D \in \mathcal{I}^{X \setminus \Lambda}$  and  $\Lambda \neq \emptyset$ .

*Proof.* For (i) (respectively (ii)) it suffices to apply Corollary 4.2 with the function  $g = -\delta_D$  (respectively  $g = \delta_D$ ), which takes the values  $\alpha_1 = -\infty$ ,  $\alpha_2 = 0$  (respectively  $\alpha_1 = 0$ ,  $\alpha_2 = +\infty$ ).  $\square$

Combining Corollaries 4.1 and 4.3, we derive the following result giving various characterizations of  $\mathcal{I}^\Lambda$  via the classes  $\mathcal{E}^{\delta_\Lambda}$  and  $\mathcal{E}^{-\delta_{X \setminus \Lambda}}$ .

**COROLLARY 4.4.** *For every set  $\Lambda \subset X$  and every set  $D \subset X$  such that  $D \neq \emptyset$  and  $D \neq X$ , the following equivalences hold:*

$$D \in \mathcal{I}^\Lambda \iff \delta_D \in \mathcal{E}^{\delta_\Lambda} \iff -\delta_{X \setminus D} \in \mathcal{E}^{-\delta_{X \setminus \Lambda}} \iff \delta_D \in \mathcal{E}^{-\delta_{X \setminus \Lambda}}.$$

*Proof.* The first equivalence is a consequence of Corollary 4.1, under the assumptions  $D \neq \emptyset$  and  $D \neq X$ . The equivalence  $D \in \mathcal{I}^\Lambda \iff -\delta_{X \setminus D} \in \mathcal{E}^{-\delta_{X \setminus \Lambda}}$  follows from Corollary 4.3(i) applied with  $X \setminus D$  (respectively  $X \setminus \Lambda$ ) in place of  $D$  (respectively  $\Lambda$ ). If  $\Lambda \neq X$ , the equivalence  $D \in \mathcal{I}^\Lambda \iff \delta_D \in \mathcal{E}^{-\delta_{X \setminus \Lambda}}$  is a consequence of Corollary 4.3(ii) applied with  $X \setminus \Lambda$  in place of  $\Lambda$ . If  $\Lambda = X$ , the equivalence becomes  $D \in \mathcal{I}^X \iff \delta_D \in \mathcal{E}^{-\omega_X}$ . Since  $\mathcal{I}^X = \{X\}$  and  $\mathcal{E}^{-\omega_X} = \{-\omega_X\}$ , the equivalence amounts to  $D = X \iff \delta_D = -\omega_X$ . The condition  $D = X$  is not realized by assumption, while the condition  $\delta_D = -\omega_X$  is never realized. It ensues that the equivalence trivially holds true if  $\Lambda = X$ .  $\square$

By combining Proposition 3.3(ii) and Proposition 4.6(iii), we obtain the following statement.

**PROPOSITION 4.7.** *Let  $X$  be a locally convex space. Let  $\Lambda \subset X$  and  $\xi^* \in X^*$ . If  $\Lambda \neq X$ , the following assertions are equivalent:*

- (a)  $[(\xi^*, \cdot) \leq 0] \in \mathcal{I}^\Lambda$ ;
- (b) for all  $\lambda \in \mathbb{R}$ ,  $[(\xi^*, \cdot) \leq \lambda] \in \mathcal{I}^\Lambda$ ;
- (c)  $(\xi^*, \cdot) \in \mathcal{E}^{-\delta_{X \setminus \Lambda}}$ ;
- (d)  $\xi^* \in -\text{dom } \sigma_{X \setminus \Lambda}$ .

*Proof.* For every  $\lambda \in \mathbb{R}$ , the set  $[(\xi^*, \cdot) \leq \lambda]$  is translated from the set  $[(\xi^*, \cdot) \leq 0]$ . Since the class  $\mathcal{I}^\Lambda$  is stable under translations, the equivalence (a)  $\iff$  (b) follows immediately. Recalling that  $\Lambda \neq X$ , we have  $\emptyset \in \mathcal{I}^\Lambda$ , hence the inclusion in (b) holds for  $\lambda = -\infty$ . Since  $X \in \mathcal{I}^\Lambda$ , the inclusion in (b) is also satisfied for  $\lambda = +\infty$ . The equivalence (b)  $\iff$  (c) can then be deduced from Proposition 4.6(iii) applied with  $X \setminus \Lambda$  in place of  $\Lambda$ . Finally, the equivalence (c)  $\iff$  (d) follows from Proposition 3.3(ii) applied with  $\varphi = -\delta_{X \setminus \Lambda}$ .  $\square$

Let us denote by  $\mathcal{C}(X)$  the class of non-empty closed convex sets of  $X$ .

**THEOREM 4.1.** *Let  $X$  be a locally convex space. Let  $\Lambda \subset X$  be such that  $\Lambda \neq X$ . For every cone  $Q \subset X^*$ , the following equivalence holds true:*

$$\{C \in \mathcal{C}(X), \text{dom } \sigma_C \subset Q\} \subset \mathcal{I}^\Lambda \iff Q \subset -\text{dom } \sigma_{X \setminus \Lambda}.$$

*Proof.* Let  $Q \subset X^*$  be a cone and assume that

$$\{C \in \mathcal{C}(X), \text{dom } \sigma_C \subset Q\} \subset \mathcal{I}^\Lambda. \tag{30}$$

Let  $\xi^* \in Q$ . Setting  $C = [(\xi^*, \cdot) \leq 0] \in \mathcal{C}(X)$ , we have  $\sigma_C = \delta_{\mathbb{R}_+ \xi^*}$ , and hence  $\text{dom } \sigma_C = \mathbb{R}_+ \xi^* \subset Q$ . In view of (30), it ensues that  $C \in \mathcal{I}^\Lambda$ . We then deduce from Proposition 4.7 that  $\xi^* \in -\text{dom } \sigma_{X \setminus \Lambda}$ . Since this is true for every  $\xi^* \in Q$ , we conclude that  $Q \subset -\text{dom } \sigma_{X \setminus \Lambda}$ .

Now assume that  $Q \subset -\text{dom } \sigma_{X \setminus \Lambda}$  and let  $C \in \mathcal{C}(X)$  be such that  $\text{dom } \sigma_C \subset Q$ . Then  $\delta_C \in \Gamma_0(X)$  with  $\text{dom } \delta_C^* \subset Q$ , and since

$$Q \subset -\text{dom } \sigma_{X \setminus \Lambda} = -\text{dom } \delta_{X \setminus \Lambda}^* = -\text{dom}(-(-\delta_{X \setminus \Lambda}))^*,$$

by Theorem 3.1 we have  $\delta_C \in \mathcal{E}^{-\delta_{X \setminus \Lambda}}$  (keep in mind  $-\delta_{X \setminus \Lambda} \neq -\omega_X$  since  $\Lambda \neq X$ ). Corollary 4.3(ii) yields that  $C \in \mathcal{I}^\Lambda$  as desired. Finally, we have shown the inclusion (30), which ends the proof.  $\square$

Applying Theorem 4.1 with  $Q = X^*$ , we immediately obtain the following result.

**COROLLARY 4.5.** *Let  $X$  be a locally convex space. Let  $\Lambda \subset X$  be such that  $\Lambda \neq X$ . Then, the following equivalence holds true:*

$$\mathcal{C}(X) \subset \mathcal{I}^\Lambda \iff \text{dom } \sigma_{X \setminus \Lambda} = X^*.$$

§5. A preorder relation on  $\mathcal{F}(X, \overline{\mathbb{R}})$  based on  $\varphi$ -envelopes. Let  $X$  be a vector space and let  $\mathcal{F}(X, \overline{\mathbb{R}})$  be the set of extended real-valued functions on  $X$ . We define the relation  $\sim$  on the space  $\mathcal{F}(X, \overline{\mathbb{R}})$  as follows: for every  $\varphi, \psi : X \rightarrow \overline{\mathbb{R}}$ ,

$$\begin{aligned} \psi \sim \varphi &\iff \text{there exist } \xi \in X \text{ and } \alpha \in \mathbb{R} \text{ such that } \psi = \varphi(\cdot - \xi) + \alpha \\ &\iff \psi \text{ is a } \varphi\text{-elementary function.} \end{aligned}$$

Clearly, the relation  $\sim$  is reflexive, symmetric and transitive, hence defines an equivalence relation. The objective of this section is to determine suitable† subsets  $\mathcal{G}$  of  $\mathcal{F}(X, \overline{\mathbb{R}})$  such that the following implication holds true for every  $\varphi, \psi \in \mathcal{G}$ :

$$\psi \in \mathcal{E}^\varphi \quad \text{and} \quad \varphi \in \mathcal{E}^\psi \implies \psi \sim \varphi. \tag{31}$$

5.1. The coercive case. For any function  $\varphi : X \rightarrow \overline{\mathbb{R}}$ , the deconvolution function  $\varphi \ominus \varphi$  defined by  $(\varphi \ominus \varphi)(x) = \sup_{y-z=x} (\varphi(y) \dot{-} \varphi(z))$  can be expressed as a  $\varphi$ -envelope via the equality  $\varphi \ominus \varphi = (\varphi_-)^\varphi$ . The next lemma shows that this function is subadditive. Recall that a function  $f : X \rightarrow \overline{\mathbb{R}}$  is said to be subadditive if for any  $x, y \in X$ ,

$$f(x + y) \leq f(x) \dot{+} f(y).$$

LEMMA 5.1. Let  $X$  be a vector space and let  $f, \varphi : X \rightarrow \overline{\mathbb{R}}$ . For any  $x, x' \in X$ , we have

$$f^\varphi(x') \leq (\varphi \ominus \varphi)(x' - x) \dot{+} f^\varphi(x).$$

Moreover, the function  $\varphi \ominus \varphi$  is subadditive.

Proof. Fix  $x, x' \in X$ . It is immediate to check that for every  $y \in X$ ,

$$\varphi(x' - y) \dot{-} f(y) \leq [\varphi(x' - y) \dot{-} \varphi(x - y)] \dot{+} [\varphi(x - y) \dot{-} f(y)].$$

Taking the supremum over  $y \in X$  and using [21, Proposition 4.a] we deduce that

$$\begin{aligned} f^\varphi(x') &\leq \sup_{y \in X} [\varphi(x' - y) \dot{-} \varphi(x - y)] \dot{+} \sup_{y \in X} [\varphi(x - y) \dot{-} f(y)], \\ &= (\varphi \ominus \varphi)(x' - x) \dot{+} f^\varphi(x), \end{aligned}$$

which yields the desired inequality. Further taking  $f = \varphi_-$  in the above inequality and using the identity  $(\varphi_-)^\varphi = \varphi \ominus \varphi$ , we obtain

$$(\varphi \ominus \varphi)(x') \leq (\varphi \ominus \varphi)(x' - x) \dot{+} (\varphi \ominus \varphi)(x),$$

hence the function  $\varphi \ominus \varphi$  is subadditive. □

† The implication (31) is not true for all  $\varphi, \psi \in \mathcal{F}(X, \overline{\mathbb{R}})$ , see a counterexample in §5.3.

If the space  $(X, \|\cdot\|)$  is normed and if the function  $\varphi$  satisfies the coercivity property  $\lim_{\|x\| \rightarrow +\infty} \varphi(x)/\|x\| = +\infty$ , the following lemma shows that  $\varphi \ominus \varphi = +\infty$  on  $X \setminus \{0\}$ .

LEMMA 5.2. *Let  $(X, \|\cdot\|)$  be a normed space and let  $\varphi : X \rightarrow \overline{\mathbb{R}}$  be an extended real-valued function. Assume that  $\varphi \neq +\omega_X$  and  $\lim_{\|x\| \rightarrow +\infty} \varphi(x)/\|x\| = +\infty$ . Then we have  $\varphi \ominus \varphi = +\infty$  on  $X \setminus \{0\}$ .*

*Proof.* Let us argue by contradiction and assume that there exists  $u \neq 0$  such that  $(\varphi \ominus \varphi)(u) < +\infty$ . Let us fix  $\bar{x} \in \text{dom } \varphi$  and observe that for every  $\dagger n \in \mathbb{N}$ ,

$$\begin{aligned} \varphi(\bar{x} + nu) \dot{-} \varphi(\bar{x}) &\leq (\varphi \ominus \varphi)(nu) \\ &\leq n(\varphi \ominus \varphi)(u) \quad \text{since } \varphi \ominus \varphi \text{ is subadditive.} \end{aligned}$$

It ensues that

$$\frac{1}{n}\varphi(\bar{x} + nu) \leq \frac{1}{n}\varphi(\bar{x}) + (\varphi \ominus \varphi)(u),$$

and taking the upper limit as  $n \rightarrow +\infty$ , we deduce that

$$\limsup_{n \rightarrow +\infty} \frac{1}{n}\varphi(\bar{x} + nu) \leq (\varphi \ominus \varphi)(u),$$

which contradicts the fact that  $\lim_{\|x\| \rightarrow +\infty} \varphi(x)/\|x\| = +\infty$ . Finally, we obtain that  $\varphi \ominus \varphi = +\infty$  on  $X \setminus \{0\}$ . □

THEOREM 5.1. *Let  $X$  be a vector space and let  $\varphi, \psi : X \rightarrow \overline{\mathbb{R}}$  be such that  $\psi \in \mathcal{E}^\varphi$  and  $\varphi \in \mathcal{E}^\psi$ .*

- (i) *If  $\varphi \ominus \varphi = +\infty$  on  $X \setminus \{0\}$ , then we have  $\psi \sim \varphi$ .*
- (ii) *Assume that  $(X, \|\cdot\|)$  is a normed space. If  $\lim_{\|x\| \rightarrow +\infty} \varphi(x)/\|x\| = +\infty$  (respectively  $\lim_{\|x\| \rightarrow +\infty} \varphi(x)/\|x\| = -\infty$ ), then we have  $\psi \sim \varphi$ .*

*Proof.* If  $\varphi = \pm\omega_X$ , it is immediate to check that  $\psi = \varphi$ . From now on, let us assume that  $\varphi \neq \pm\omega_X$ . Since  $\psi \in \mathcal{E}^\varphi$  and  $\varphi \in \mathcal{E}^\psi$ , there exist  $f, g : X \rightarrow \overline{\mathbb{R}}$  such that  $-\psi = (-\varphi) \nabla f$  and  $-\varphi = (-\psi) \nabla g$ . It ensues that

$$-\varphi = (-\varphi) \nabla (f \nabla g). \tag{32}$$

Now observe that

$$\begin{aligned} (-\varphi) \nabla (f \nabla g) \geq -\varphi &\iff (-\varphi)(x - y) \dot{+} (f \nabla g)(y) \geq -\varphi(x) \\ &\quad \text{for all } x, y \in X \\ &\iff (f \nabla g)(y) \geq \varphi(x - y) \dot{-} \varphi(x) \quad \text{for all } x, y \in X \\ &\iff (f \nabla g)(y) \geq \sup_{x \in X} (\varphi(x - y) \dot{-} \varphi(x)) \\ &\quad \text{for all } y \in X \\ &\iff f \nabla g \geq [\varphi \ominus \varphi]_-. \end{aligned}$$

$\dagger$  Here  $\mathbb{N}$  denotes the set of positive integers.



(i) Assume that  $\varphi \ominus \varphi = +\infty$  on  $X \setminus \{0\}$ . We then deduce from the above inequality that

$$f \nabla g = +\infty \quad \text{on } X \setminus \{0\}. \tag{33}$$

If  $f \nabla g = \omega_X$ , we infer from (32) that  $\varphi = -\omega_X$ , thus implying in turn that  $\psi = -\omega_X$ . If  $f \nabla g \neq \omega_X$ , equality (33) shows that  $\text{dom}(f \nabla g) = \{0\}$ . Recalling that  $\text{dom}(f \nabla g) = \text{dom } f + \text{dom } g$ , we deduce that  $\text{dom } f + \text{dom } g = \{0\}$ . Hence, there exists  $\xi \in X$  such that  $\text{dom } f = \{\xi\}$  and  $\text{dom } g = \{-\xi\}$ . We infer that

$$-\psi = (-\varphi) \nabla f = (-\varphi)(\cdot - \xi) \dot{+} f(\xi) \tag{34}$$

and

$$-\varphi = (-\psi) \nabla g = (-\psi)(\cdot + \xi) \dot{+} g(-\xi). \tag{35}$$

If  $f(\xi) \in \mathbb{R}$ , we obtain from (34) that  $\psi = \varphi(\cdot - \xi) - f(\xi)$  and therefore  $\psi \sim \varphi$ . If  $g(-\xi) \in \mathbb{R}$ , equality (35) shows that  $\varphi = \psi(\cdot + \xi) - g(-\xi)$ , and hence  $\varphi \sim \psi$ . On the other hand, if  $f(\xi) = g(-\xi) = -\infty$ , we deduce from (34)–(35) that

$$-\psi \leq (-\varphi)(\cdot - \xi) \quad \text{and} \quad -\varphi \leq (-\psi)(\cdot + \xi),$$

thus implying that  $\psi = \varphi(\cdot - \xi)$  and therefore  $\psi \sim \varphi$ .

(ii) First assume that  $\lim_{\|x\| \rightarrow +\infty} \varphi(x)/\|x\| = +\infty$ . We infer from Lemma 5.2 that  $\varphi \ominus \varphi = +\infty$  on  $X \setminus \{0\}$  and we conclude with (i).

Now assume that  $\lim_{\|x\| \rightarrow +\infty} \varphi(x)/\|x\| = -\infty$ . From Lemma 5.2, we deduce that  $(-\varphi) \ominus (-\varphi) = +\infty$  on  $X \setminus \{0\}$ . Recalling that

$$(-\varphi) \ominus (-\varphi) = (-\varphi_-)^{-\varphi} = \varphi^{\varphi^-} = [(\varphi_-)^{\varphi}]_- = [\varphi \ominus \varphi]_-,$$

we infer that  $\varphi \ominus \varphi = +\infty$  on  $X \setminus \{0\}$  and we conclude again with (i). □

Let us define the relation  $\leq$  on  $\mathcal{F}(X, \overline{\mathbb{R}})$  by

$$\psi \leq \varphi \iff \psi \in \mathcal{E}^\varphi.$$

The relation  $\leq$  is clearly reflexive, and also transitive in view of Proposition 3.2(iii). It is compatible with the equivalence relation  $\sim$ , i.e.

$$\varphi \sim \varphi', \quad \psi \sim \psi' \quad \text{and} \quad \psi \leq \varphi \implies \psi' \leq \varphi'.$$

It ensues that we can properly define the relation  $\leq$  on the quotient set  $\mathcal{F}(X, \overline{\mathbb{R}})/\sim$ . The relation  $\leq$  so defined on  $\mathcal{F}(X, \overline{\mathbb{R}})/\sim$  is clearly reflexive and transitive, hence it is a preorder. Let us denote by  $\mathcal{G}$ ,  $\mathcal{G}'$  and  $\mathcal{G}''$  the following respective sets:

$$\begin{aligned} \mathcal{G} &= \{f : X \rightarrow \overline{\mathbb{R}}, f \ominus f = +\infty \text{ on } X \setminus \{0\}\}, \\ \mathcal{G}' &= \left\{f : X \rightarrow \overline{\mathbb{R}}, \lim_{\|x\| \rightarrow +\infty} f(x)/\|x\| = +\infty\right\}, \\ \mathcal{G}'' &= \left\{f : X \rightarrow \overline{\mathbb{R}}, \lim_{\|x\| \rightarrow +\infty} f(x)/\|x\| = -\infty\right\}. \end{aligned}$$

Theorem 5.1 expresses that for every  $\varphi, \psi \in \mathcal{G}$  (respectively  $\mathcal{G}', \mathcal{G}''$ ), we have

$$\psi \preceq \varphi, \quad \varphi \preceq \psi \implies \psi \sim \varphi.$$

Hence, the induced relation  $\preceq$  on the quotient set  $\mathcal{G}/\sim$  (respectively  $\mathcal{G}'/\sim, \mathcal{G}''/\sim$ ) is antisymmetric, thus giving rise to an order relation.

Let us now specialize the result of Theorem 5.1 in the case of sets. We define the equivalence relation  $\sim$  on  $\mathcal{P}(X)$  by

$$C \sim D \iff \text{there exists } \xi \in X \text{ such that } D = C + \xi,$$

along with the preorder relation  $\preceq$  on  $\mathcal{P}(X)$  by

$$C \preceq D \iff C \in \mathcal{I}^D.$$

Recall that the star-difference  $C \overset{*}{\ominus} C$  is defined by

$$C \overset{*}{\ominus} C = \bigcap_{x \in C} C - x = (-C)^C.$$

By applying Theorem 5.1 with indicator functions, we obtain the following corollary.

**COROLLARY 5.1.** *Let  $X$  be a vector space and let  $\Gamma, \Delta \subset X$  be such that  $\Delta \in \mathcal{I}^\Gamma$  and  $\Gamma \in \mathcal{I}^\Delta$ .*

- (i) *If  $\Gamma \overset{*}{\ominus} \Gamma = \{0\}$ , then we have  $\Delta \sim \Gamma$ .*
- (ii) *Assume that  $(X, \|\cdot\|)$  is a normed space. If the set  $\Gamma$  (respectively  $X \setminus \Gamma$ ) is bounded, then we have  $\Delta \sim \Gamma$ .*

*Proof.* If  $\Gamma \in \{\emptyset, X\}$  (respectively  $\Delta \in \{\emptyset, X\}$ ), it is immediate to check that  $\Delta = \Gamma$ . Let us now assume that  $\Gamma \notin \{\emptyset, X\}$  and  $\Delta \notin \{\emptyset, X\}$ . In view of Corollary 4.1, the assumptions  $\Delta \in \mathcal{I}^\Gamma$  and  $\Gamma \in \mathcal{I}^\Delta$  imply that  $\delta_\Delta \in \mathcal{E}^{\delta_\Gamma}$  and  $\delta_\Gamma \in \mathcal{E}^{\delta_\Delta}$ .

- (i) Assume that  $\Gamma \overset{*}{\ominus} \Gamma = \{0\}$ . Then, by (5) and Proposition 4.4(i), we have

$$\delta_\Gamma \ominus \delta_\Gamma = (\delta_{-\Gamma})^{\delta_\Gamma} = \delta_{(-\Gamma)\Gamma} = \delta_{\Gamma \overset{*}{\ominus} \Gamma} = \delta_{\{0\}}.$$

By applying Theorem 5.1(i) with  $\varphi = \delta_\Gamma$  and  $\psi = \delta_\Delta$ , we obtain that  $\delta_\Delta \sim \delta_\Gamma$  and hence  $\Delta \sim \Gamma$ .

(ii) First assume that  $\Gamma$  is bounded. Then the indicator function  $\delta_\Gamma$  is coercive and we deduce from Lemma 5.2 that  $\delta_\Gamma \ominus \delta_\Gamma = +\infty$  on  $X \setminus \{0\}$ . This implies that  $\Gamma \overset{*}{\ominus} \Gamma = \{0\}$  and we conclude with (i). Now assume that  $X \setminus \Gamma$  is bounded. From what precedes, we have  $(X \setminus \Gamma) \overset{*}{\ominus} (X \setminus \Gamma) = \{0\}$ . Observing that

$$\Gamma \overset{*}{\ominus} \Gamma = (-\Gamma)^\Gamma = (X \setminus \Gamma)^{-X \setminus \Gamma} = -[(-X \setminus \Gamma)^{X \setminus \Gamma}] = -[(X \setminus \Gamma) \overset{*}{\ominus} (X \setminus \Gamma)],$$

we infer that  $\Gamma \overset{*}{\ominus} \Gamma = \{0\}$  and we conclude again with (i). □

Let us denote by  $\mathcal{Q}$ ,  $\mathcal{Q}'$  and  $\mathcal{Q}''$  the following respective sets:

$$\begin{aligned} \mathcal{Q} &= \{C \subset X, C \overset{*}{=} \{0\}\}, \\ \mathcal{Q}' &= \{C \subset X, C \text{ is bounded}\}, \\ \mathcal{Q}'' &= \{C \subset X, X \setminus C \text{ is bounded}\}. \end{aligned}$$

The above corollary expresses that for every  $\Gamma, \Delta \in \mathcal{Q}$  (respectively  $\mathcal{Q}', \mathcal{Q}''$ ), we have

$$\Delta \preceq \Gamma, \quad \Gamma \preceq \Delta \implies \Delta \sim \Gamma.$$

Hence, the induced relation  $\preceq$  on the quotient set  $\mathcal{Q}/\sim$  (respectively  $\mathcal{Q}'/\sim, \mathcal{Q}''/\sim$ ) is antisymmetric, thus giving rise to an order relation.

5.2. *The case  $\varphi, \psi \in -\Gamma_0(X)$ .* Let us first state a result that will be a key ingredient for the next theorem.

LEMMA 5.3. *Let  $X$  be a vector space, let  $D \subset X$  be a convex set and let us denote by  $\text{Aff}(D)$  the affine space generated by  $D$ . Assume that a real-valued function  $h : D \rightarrow \mathbb{R}$  is both convex and concave. Then there exists a unique affine function  $\tilde{h} : \text{Aff}(D) \rightarrow \mathbb{R}$  such that  $\tilde{h}|_D = h$ .*

For a proof of this result, the reader is referred to [35]. By extending affinely the function  $\tilde{h}$  to the whole space  $X$ , we deduce from the above result that there exists a linear function  $\ell : X \rightarrow \mathbb{R}$  along with  $\alpha \in \mathbb{R}$  such that  $h = \ell|_D + \alpha$ .

In view of stating the next theorem, given a locally convex space  $X$  recall that the Mackey topology  $\tau(X^*, X)$  on  $X^*$  is defined as the finest locally convex topology  $\mathcal{T}$  on  $X^*$  such that the topological dual of  $(X^*, \mathcal{T})$  coincides with  $X$ . If  $(X, \|\cdot\|)$  is normed, this topology is exactly that associated with the dual norm  $\|\cdot\|_{X^*}$  provided that  $(X, \|\cdot\|)$  is a reflexive Banach space.

THEOREM 5.2. *Let  $X$  be a locally convex space. Let  $\varphi, \psi : X \rightarrow \overline{\mathbb{R}}$  be functions such that  $\psi \in \mathcal{E}^\varphi$  and  $\varphi \in \mathcal{E}^\psi$ . Assume that either:*

- *the space  $X$  is finite-dimensional; or*
- *one of the functions  $(-\varphi)^*$  and  $(-\psi)^*$  is  $\tau(X^*, X)$ -continuous at some point and finite at this point.*

*Then we have  $(-\varphi)^{**} \sim (-\psi)^{**}$ . If each of the functions  $-\varphi$  and  $-\psi$  has a continuous affine minorant, then  $\overline{\text{co}}(-\varphi) \sim \overline{\text{co}}(-\psi)$ . In particular, if  $-\varphi \in \Gamma_0(X)$  and  $-\psi \in \Gamma_0(X)$ , then we have  $\varphi \sim \psi$ .*

*Proof.* By assumption, we have  $-\psi = (-\varphi) \nabla f$  and  $-\varphi = (-\psi) \nabla g$ , for some  $f, g : X \rightarrow \overline{\mathbb{R}}$ . Taking the conjugates, we obtain that

$$(-\psi)^* = (-\varphi)^* \dot{+} f^* \quad \text{and} \quad (-\varphi)^* = (-\psi)^* \dot{+} g^*. \tag{36}$$

First observe that if one of the functions  $(-\varphi)^*, (-\psi)^*, f^*$  or  $g^*$  is equal to  $-\omega_{X^*}$ , then equalities (36) imply that  $(-\varphi)^* = (-\psi)^* = -\omega_{X^*}$ . This implies

in turn that  $\varphi = \psi = -\omega_X$  and the conclusion is satisfied. From now on, let us assume that the functions  $(-\varphi)^*$ ,  $(-\psi)^*$ ,  $f^*$  and  $g^*$  differ from  $-\omega_{X^*}$ . From the first equality of (36), we deduce that  $\text{dom}(-\psi)^* \subset \text{dom}(-\varphi)^*$ , while the second equality of (36) yields  $\text{dom}(-\varphi)^* \subset \text{dom}(-\psi)^*$ . Finally, the domains of  $(-\varphi)^*$  and  $(-\psi)^*$  coincide and both functions are finite on their common domain  $D$ . If the set  $D$  is empty, then  $(-\varphi)^* = (-\psi)^* = \omega_{X^*}$ . This implies that  $(-\varphi)^{**} = (-\psi)^{**} = -\omega_X$ , hence the conclusion is trivially satisfied. Without loss of generality, we now assume that  $D \neq \emptyset$ . By combining both equalities of (36), we obtain

$$(-\varphi)^* = (-\varphi)^* + f^* + g^*.$$

It ensues that  $f^* + g^* = 0$  on  $D$ . Hence, the function  $f^*|_D$  is finite-valued on  $D$  and both convex and concave. By applying the previous lemma with  $h = f^*|_D$ , we obtain that there exist a linear function  $\ell : X^* \rightarrow \mathbb{R}$  and  $\alpha \in \mathbb{R}$  such that  $f^* = \ell + \alpha$  on  $D$ . Coming back to the first equality of (36), we deduce that

$$(-\psi)^* = (-\varphi)^* + \ell + \alpha.$$

Observe that the above equality holds true on the whole space  $X^*$ , since the functions  $(-\varphi)^*$  and  $(-\psi)^*$  are equal to  $+\infty$  outside  $D$ . Taking the conjugate of each member, we find for every  $\xi \in X$ ,

$$(-\psi)^{**}(\xi) = \sup_{x^* \in X^*} [\langle x^*, \xi \rangle - (-\varphi)^*(x^*) - \ell(x^*) - \alpha]. \tag{37}$$

Let us now show that the linear function  $\ell$  is  $\tau(X^*, X)$ -continuous on  $X^*$ .

LEMMA 5.1. *Under the assumptions of Theorem 5.2, the function  $\ell : X^* \rightarrow \mathbb{R}$  is  $\tau(X^*, X)$ -continuous on  $X^*$ .*

*Proof of Lemma 5.1.* If the space  $X$  is finite-dimensional, the assertion is obvious. Now assume that the function  $(-\varphi)^*$  is  $\tau(X^*, X)$ -continuous at some  $\bar{x}^* \in X^*$  and finite at this point. There exist a  $\tau(X^*, X)$ -neighborhood  $W$  of  $\bar{x}^*$  and  $M \in \mathbb{R}$  such that  $(-\varphi)^* \leq M$  on  $W$ . We deduce from (37) that for every  $\xi \in X$ ,

$$\begin{aligned} (-\psi)^{**}(\xi) &\geq \sup_{x^* \in W} [\langle x^*, \xi \rangle - (-\varphi)^*(x^*) - \ell(x^*) - \alpha] \\ &\geq \sup_{x^* \in W} [\langle x^*, \xi \rangle - \ell(x^*)] - M - \alpha. \end{aligned}$$

Let us argue by contradiction and assume that  $\ell$  is not  $\tau(X^*, X)$ -continuous on  $X^*$ . Since the linear function  $\langle \cdot, \xi \rangle - \ell$  is not  $\tau(X^*, X)$ -continuous on  $X^*$ , the above supremum equals  $+\infty$ . It ensues that  $(-\psi)^{**} = \omega_X$ , and hence  $-\psi = \omega_X$ . Recalling that  $-\varphi = (-\psi) \nabla g$ , we deduce that  $-\varphi = \omega_X$ . This implies in turn that  $(-\varphi)^* = -\omega_{X^*}$ , a contradiction with  $(-\varphi)^*(\bar{x}^*) \in \mathbb{R}$ . We conclude that the linear function  $\ell$  is  $\tau(X^*, X)$ -continuous on  $X^*$ . Since  $\varphi$  and  $\psi$  play symmetric roles, the same conclusion holds true if the function  $(-\psi)^*$  is assumed to be  $\tau(X^*, X)$ -continuous at some  $\tilde{x}^* \in X^*$  and finite at this point. □

From the previous lemma and the definition of the Mackey topology  $\tau(X^*, X)$ , there exists  $x \in X$  such that  $\ell(x^*) = \langle x^*, x \rangle$  for every  $x^* \in X^*$ . In view of (37), we deduce that

$$(-\psi)^{**}(\xi) = \sup_{x^* \in X^*} [\langle x^*, \xi - x \rangle - (-\varphi)^*(x^*)] - \alpha = (-\varphi)^{**}(\xi - x) - \alpha.$$

Since this is true for every  $\xi \in X$ , we conclude that  $(-\psi)^{**} \sim (-\varphi)^{**}$ . If the function  $(-\varphi)$  (respectively  $(-\psi)$ ) admits a continuous affine minorant, we have  $(-\varphi)^{**} = \overline{\text{co}}(-\varphi)$  (respectively  $(-\psi)^{**} = \overline{\text{co}}(-\psi)$ ). We infer that  $\overline{\text{co}}(-\psi) \sim \overline{\text{co}}(-\varphi)$ . The last assertion of the statement is a direct consequence of what precedes.  $\square$

*Remark 5.1.* If the normed space  $(X, \|\cdot\|)$  is reflexive, the  $\tau(X^*, X)$ -continuity assumption on  $(-\varphi)^*$  (respectively  $(-\psi)^*$ ) amounts to the continuity assumption with respect to the dual norm  $\|\cdot\|_{X^*}$ .

Theorem 5.2 implies that the relation  $\preceq$  defines an order on the following set:

$$\{\varphi \in -\Gamma_0(X), (-\varphi)^* \text{ is } \tau(X^*, X)\text{-continuous at some point}\} / \sim.$$

If the space  $X$  is finite-dimensional, the relation  $\preceq$  is an order on the set  $(-\Gamma_0(X)) / \sim$ .

By applying Theorem 5.2 with the opposite of indicator functions, we obtain the following corollary.

**COROLLARY 5.2.** *Let  $X$  be a locally convex space. Let  $\Gamma, \Delta \subset X$  be such that  $\Delta \in \mathcal{I}^\Gamma$  and  $\Gamma \in \mathcal{I}^\Delta$ . Assume that either:*

- *the space  $X$  is finite-dimensional; or*
- *one of the functions  $\sigma_{X \setminus \Gamma}$  and  $\sigma_{X \setminus \Delta}$  is  $\tau(X^*, X)$ -continuous at some point.*

*Then we have  $\overline{\text{co}}(X \setminus \Gamma) \sim \overline{\text{co}}(X \setminus \Delta)$ . In particular, if the sets  $X \setminus \Gamma$  and  $X \setminus \Delta$  are closed and convex, then  $\Gamma \sim \Delta$ .*

*Proof.* From Corollary 4.3(i), condition  $\Delta \in \mathcal{I}^\Gamma$  (respectively  $\Gamma \in \mathcal{I}^\Delta$ ) is equivalent to  $-\delta_{X \setminus \Delta} \in \mathcal{E}^{-\delta_{X \setminus \Gamma}}$  (respectively  $-\delta_{X \setminus \Gamma} \in \mathcal{E}^{-\delta_{X \setminus \Delta}}$ ). By applying Theorem 5.2 with  $\varphi = -\delta_{X \setminus \Gamma}$  and  $\psi = -\delta_{X \setminus \Delta}$ , we obtain the existence of  $\xi \in X$  and  $\alpha \in \mathbb{R}$  such that

$$\overline{\text{co}}(\delta_{X \setminus \Delta}) = [\overline{\text{co}}(\delta_{X \setminus \Gamma})](\cdot - \xi) - \alpha.$$

We immediately deduce that  $\overline{\text{co}}(X \setminus \Delta) = \overline{\text{co}}(X \setminus \Gamma) + \xi$ . The last assertion of the statement is a direct consequence of what precedes.  $\square$

5.3. *A counterexample.* Let us start with a preliminary result.

**LEMMA 5.4.** *Let  $X$  be a topological vector space and let  $G$  be a dense additive subgroup of  $X$ . Assume that  $K \subset X$  is an open set such that  $K + K \subset K$  and  $0 \in \text{cl}(K)$ . Then we have:*

(i) for all  $\xi, \xi' \in X$ ,

$$[G \cap (K + \xi)] + [G \cap (K + \xi')] = G \cap (K + \xi + \xi');$$

(ii) if, in addition,  $\text{cl}(K) \cap -\text{cl}(K) = \{0\}$ , then

$$G \cap (K + \xi) = (G \cap K) + \xi' \implies \xi = \xi';$$

if  $G \neq X$  and  $\xi \in X \setminus G$ , there is no  $\xi' \in X$  such that  $G \cap (K + \xi) = (G \cap K) + \xi'$ .

*Proof.* (i) Let us fix  $\xi, \xi' \in X$  and let us prove the inclusion from the left to the right. Observe that

$$[G \cap (K + \xi)] + [G \cap (K + \xi')] \subset G + G$$

and

$$[G \cap (K + \xi)] + [G \cap (K + \xi')] \subset (K + \xi) + (K + \xi').$$

Since  $G + G \subset G$  and  $K + K \subset K$ , we deduce that

$$[G \cap (K + \xi)] + [G \cap (K + \xi')] \subset G \cap (K + \xi + \xi').$$

Now let us establish the reverse inclusion. Let  $x \in G \cap (K + \xi + \xi')$ . Observe that the open set  $K + \xi + \xi' - x$  contains 0. Recalling that  $0 \in \text{cl}(K)$ , we have  $(K + \xi + \xi' - x) \cap -K \neq \emptyset$ , hence  $(K + \xi - x) \cap (-K - \xi') \neq \emptyset$ . Since the set  $K$  is open, the set  $(K + \xi - x) \cap (-K - \xi')$  is open. By using the density of  $G$  in  $X$ , we deduce that

$$G \cap (K + \xi - x) \cap (-K - \xi') \neq \emptyset.$$

Since  $G = -G$ , the above property can be rewritten as

$$[G \cap (K + \xi - x)] \cap [-G \cap (-K - \xi')] \neq \emptyset,$$

which is, in turn, equivalent to

$$0 \in [G \cap (K + \xi - x)] + [G \cap (K + \xi')].$$

Recalling that  $x \in G$ , we have  $G = G - x$ , hence

$$G \cap (K + \xi - x) = [G \cap (K + \xi)] - x.$$

In view of the latter inclusion, we conclude that

$$x \in [G \cap (K + \xi)] + [G \cap (K + \xi')].$$

The inclusion

$$G \cap (K + \xi + \xi') \subset [G \cap (K + \xi)] + [G \cap (K + \xi')]$$

is proved.

(ii) Let us assume that  $G \cap (K + \xi) = (G \cap K) + \xi'$  for some  $\xi, \xi' \in X$ . We deduce that  $G \cap (K + \xi) \subset K + \xi'$ . By using the openness of the set  $K + \xi$  along with the density of  $G$  in  $X$ , we easily infer that  $K + \xi \subset \text{cl}(K) + \xi'$ . This implies, in turn, that  $\text{cl}(K) + \xi \subset \text{cl}(K) + \xi'$  and since  $0 \in \text{cl}(K)$ , we obtain  $\xi - \xi' \in \text{cl}(K)$ . By a symmetric argument, we find  $\xi' - \xi \in \text{cl}(K)$ , hence  $\xi - \xi' \in \text{cl}(K) \cap -\text{cl}(K)$ . Since  $\text{cl}(K) \cap -\text{cl}(K) = \{0\}$  by assumption, we conclude that  $\xi = \xi'$ .

Now let  $\xi \in X \setminus G$  and assume that there exists  $\xi' \in X$  such that  $G \cap (K + \xi) = (G \cap K) + \xi'$ . From what precedes, we have  $\xi' = \xi$  and, hence,  $G \cap (K + \xi) = (G + \xi) \cap (K + \xi)$ . On the other hand, the assumption  $\xi \in X \setminus G$  implies that the sets  $G$  and  $G + \xi$  are disjoint. We deduce that  $G \cap (K + \xi) = \emptyset$ , a contradiction since the non-empty set  $K$  is open and the set  $G$  is dense in  $X$ . □

Let us now build an example of sets  $\Gamma, \Delta \subset X$  satisfying  $\Delta \in \mathcal{I}^\Gamma$  and  $\Gamma \in \mathcal{I}^\Delta$ , but with  $\Delta$  and  $\Gamma$  not equal up to a translation. We are given an open set  $K \subset X$  such that  $K + K \subset K$  and  $\text{cl}(K) \cap -\text{cl}(K) = \{0\}$ , along with a dense additive subgroup  $G \subset X$  such that  $G \neq X$ . Define the sets  $C, U, V \subset X$  respectively by

$$C = G \cap K; \quad U = G \cap (K + \xi); \quad V = G \cap (K - \xi),$$

where  $\xi \in X \setminus G$ . In view of Lemma 5.4(i), the set  $D = C + U$  satisfies

$$D = G \cap (K + \xi) \quad \text{and} \quad D + V = G \cap K = C.$$

Lemma 5.4(ii) shows that the set  $D$  is not translated from  $C$ . Defining the complementary sets  $\Gamma = X \setminus C$  and  $\Delta = X \setminus D$ , we have

$$\Delta = X \setminus (C + U) = U^{X \setminus C} = U^\Gamma \in \mathcal{I}^\Gamma \tag{38}$$

and

$$\Gamma = X \setminus (D + V) = V^{X \setminus D} = V^\Delta \in \mathcal{I}^\Delta. \tag{39}$$

From what precedes, the set  $\Delta$  is not translated from  $\Gamma$ . The above counterexample for sets obviously furnishes a counterexample for functions. Indeed, we deduce from (38)–(39) that the indicator functions  $\delta_\Gamma$  and  $\delta_\Delta$  satisfy  $\delta_\Delta \in \mathcal{E}^{\delta_\Gamma}$  and  $\delta_\Gamma \in \mathcal{E}^{\delta_\Delta}$ , but the functions  $\delta_\Gamma$  and  $\delta_\Delta$  are not equal up to a translation.

By particularizing the above sets  $G, K \subset X$ , one obtains various counterexamples. If  $X = \mathbb{R}$ , one can take  $G = \mathbb{Q}, K = ]0, +\infty[$  and  $\xi \in \mathbb{R} \setminus \mathbb{Q}$ . On the other hand, if  $X$  is infinite-dimensional, one can assume that  $G$  is a dense subspace of  $X$  and that  $K$  is an open convex cone such that  $\text{cl}(K)$  is pointed. This furnishes a counterexample with convex sets  $C, D \subset X$ .

§6. *Cases of either superadditivity or subadditivity of  $\varphi$ .* Let us first recall that a function  $\varphi : X \rightarrow \overline{\mathbb{R}}$  is said to be superadditive (respectively subadditive) if for all  $x, y \in X$ ,

$$\varphi(x + y) \geq \varphi(x) + \varphi(y) \quad (\text{respectively } \varphi(x + y) \leq \varphi(x) + \varphi(y)).$$

Let us start with a preliminary result.

LEMMA 6.1. *Let  $X$  be a vector space. Let  $h, k : X \rightarrow \overline{\mathbb{R}}$  and assume that  $k(0) = 0$ . Then we have*

$$\begin{aligned}
 h = h \Delta k &\iff h(x) \geq h(y) \dot{+} k(x - y) \quad \text{for all } x, y \in X \\
 &\iff h(y) \leq h(x) \dot{+} (-k_-)(y - x) \quad \text{for all } x, y \in X \\
 &\iff h = h \nabla (-k_-).
 \end{aligned}$$

*As a consequence, the function  $k$  is superadditive if and only if  $k = k \Delta k$ , which is, in turn, equivalent to  $k = k \nabla (-k_-)$ .*

*Proof.* If  $h = h \Delta k$ , then the definition of  $h \Delta k$  entails that  $h(x) \geq h(y) \dot{+} k(x - y)$  for all  $x, y \in X$ . Conversely, if this inequality holds true for every  $x, y \in X$ , we have

$$h(x) \geq \sup_{y \in X} h(y) \dot{+} k(x - y) \geq h(x) + k(0) = h(x),$$

for every  $x \in X$ . This implies that  $h(x) = (h \Delta k)(x)$  for every  $x \in X$  and the first equivalence is proved.

For the second equivalence, observe that for all  $x, y \in X$ ,

$$\begin{aligned}
 h(x) \geq h(y) \dot{+} k(x - y) &\iff \\
 h(y) \leq h(x) \dot{+} (-k)(x - y) &= h(x) \dot{+} (-k_-)(y - x).
 \end{aligned}$$

The proof of the third equivalence follows the same lines as the first one. For the last assertion, observe that  $k$  is superadditive if and only if  $k(x) \geq k(y) \dot{+} k(x - y)$  for all  $x, y \in X$ . It suffices then to use what precedes with  $h = k$ . □

Through the above lemma, the following theorem provides, in particular, various characterizations of the class  $\mathcal{E}^\varphi$  when  $\varphi$  is superadditive.

THEOREM 6.1. *Let  $X$  be a vector space. Let  $\varphi : X \rightarrow \overline{\mathbb{R}}$  be a superadditive function satisfying  $\varphi(0) = 0$ .*

- (a) *For a function  $g : X \rightarrow \overline{\mathbb{R}}$ , the following assertions are equivalent:*
  - (i)  $g \in \mathcal{E}^\varphi$ ;
  - (ii)  $g = g \Delta \varphi$ ;
  - (iii)  $g(x) \geq g(y) \dot{+} \varphi(x - y)$  for all  $x, y \in X$ ;
  - (iv)  $g(y) \leq g(x) \dot{+} (-\varphi_-)(y - x)$  for all  $x, y \in X$ ;
  - (v)  $g = g \nabla (-\varphi_-)$ ;
  - (vi)  $-g \in \mathcal{E}^{\varphi_-}$ .
- (b) *For every function  $f : X \rightarrow \overline{\mathbb{R}}$ ,  $f \nabla (-\varphi_-)$  is the greatest  $\varphi$ -envelope that is majorized by  $f$ , while  $f \Delta \varphi$  is the lowest  $\varphi$ -envelope that is minorized by  $f$ .*
- (c) *The following inclusion holds true:  $\mathcal{E}^{-\varphi} \subset \mathcal{E}^{\varphi_-}$ .*



*Proof.* (a) Let us assume that  $g \in \mathcal{E}^\varphi$ . Then there exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $g = f^\varphi = (-f) \Delta \varphi$ . Using the superadditivity of  $\varphi$  and the last assertion of Lemma 6.1, we have

$$g \Delta \varphi = ((-f) \Delta \varphi) \Delta \varphi = (-f) \Delta (\varphi \Delta \varphi) = (-f) \Delta \varphi = g.$$

This shows that (i)  $\implies$  (ii). Conversely, if  $g = g \Delta \varphi$  then  $g = (-g)^\varphi$  and clearly  $g \in \mathcal{E}^\varphi$ . The equivalences (ii)  $\iff$  (iii)  $\iff$  (iv)  $\iff$  (v) follow directly from Lemma 6.1. For the equivalence (v)  $\iff$  (vi), observe that

$$g = g \nabla (-\varphi_-) \iff -g = (-g) \Delta \varphi_-,$$

and invoke the equivalence (i)  $\iff$  (ii).

(b) Let  $f : X \rightarrow \overline{\mathbb{R}}$  and take  $g = f \nabla (-\varphi_-) = -f^{\varphi_-}$ . By using the implication (vi)  $\implies$  (ii) in (a), we obtain  $g = g \Delta \varphi$ , thus implying that  $g = (f^{\varphi_-})^\varphi$ . Hence,  $f \nabla (-\varphi_-)$  coincides with  $(f^{\varphi_-})^\varphi$ , which is by property (6) the greatest element of  $\mathcal{E}^\varphi$  that is majorized by  $f$ . Replacing  $f$  (respectively  $\varphi$ ) with  $-f$  (respectively  $\varphi_-$ ) and taking the opposite, we deduce that  $f \Delta \varphi$  is the lowest element of  $-\mathcal{E}^{\varphi_-}$  that is minorized by  $f$ . It suffices then to recall that  $\mathcal{E}^{\varphi_-} = -\mathcal{E}^\varphi$ , see the equivalence (i)  $\iff$  (vi) in (a).

(c) Since  $\varphi \in \mathcal{E}^\varphi$ , we have  $-\varphi \in -\mathcal{E}^\varphi = \mathcal{E}^{\varphi_-}$ . In view of Proposition 3.2(iii), we infer that  $\mathcal{E}^{-\varphi} \subset \mathcal{E}^{\varphi_-}$ . □

*Example 6.1.* Assume that  $(X, \|\cdot\|)$  is a normed space. For  $k \geq 0$  and  $\alpha \in ]0, 1]$ , take  $\varphi = -k\|\cdot\|^\alpha$ . Observe that for all  $x, y \in X$ ,

$$\|x + y\|^\alpha \leq (\|x\| + \|y\|)^\alpha \leq \|x\|^\alpha + \|y\|^\alpha. \tag{40}$$

It ensues that the function  $\|\cdot\|^\alpha$  is subadditive, hence  $\varphi$  is superadditive. From Theorem 6.1(a), we deduce that

$$f \in \mathcal{E}^{-k\|\cdot\|^\alpha} \iff f(x) \geq f(y) - k\|x - y\|^\alpha \quad \text{for all } x, y \in X. \tag{41}$$

By reversing the roles of  $x$  and  $y$ , we immediately obtain

$$f \in \mathcal{E}^{-k\|\cdot\|^\alpha} \iff f(x) \leq f(y) + k\|x - y\|^\alpha \quad \text{for all } x, y \in X. \tag{42}$$

If  $f(y) = +\infty$  (respectively  $f(y) = -\infty$ ) for some  $y \in X$ , we deduce from (41) (respectively (42)) that  $f = \omega_X$  (respectively  $f = -\omega_X$ ). On the other hand, if the function  $f$  is finite-valued, we infer from (41)–(42) that  $|f(x) - f(y)| \leq k\|x - y\|^\alpha$  for all  $x, y \in X$ . This implies that

$$\begin{aligned} \mathcal{E}^{-k\|\cdot\|^\alpha} &= \{f : X \rightarrow \mathbb{R}, |f(x) - f(y)| \leq k\|x - y\|^\alpha \text{ for all } x, y \in X\} \cup \{\omega_X, -\omega_X\} \\ &= \{f : X \rightarrow \mathbb{R}, f \text{ is } \alpha\text{-H\"olderian with constant } k\} \cup \{\omega_X, -\omega_X\}. \end{aligned}$$

From Theorem 6.1(b), we deduce that  $f \nabla k\|\cdot\|^\alpha$  (respectively  $f \Delta (-k\|\cdot\|^\alpha)$ ) is the greatest (respectively lowest)  $\varphi$ -envelope that is majorized (respectively minorized) by  $f$ . Since the map  $\|\cdot\|^\alpha$  is even, Theorem 6.1(c) shows that  $\mathcal{E}^{k\|\cdot\|^\alpha} \subset \mathcal{E}^{-k\|\cdot\|^\alpha}$ .

Now assume that  $\alpha = 1$ . From what precedes, we obtain that

$$\mathcal{E}^{-k\|\cdot\|} = \{f : X \rightarrow \mathbb{R}, f \text{ is } k\text{-Lipschitz continuous}\} \cup \{\omega_X, -\omega_X\}.$$

The Pasch–Hausdorff regularization of  $f$ , defined by  $l_k(f) = f \nabla k\|\cdot\|$ , is the greatest function of  $\mathcal{E}^{-k\|\cdot\|}$  that is majorized by  $f$ . On the other hand,  $f \Delta (-k\|\cdot\|)$  is the lowest function of  $\mathcal{E}^{-k\|\cdot\|}$  that is minorized by  $f$ . The inclusion  $\mathcal{E}^{k\|\cdot\|} \subset \mathcal{E}^{-k\|\cdot\|}$  shows that the  $k\|\cdot\|$ -envelopes are either  $k$ -Lipschitz continuous or equal to  $\pm\omega_X$ . The convexity of  $\|\cdot\|$  implies that  $k\|\cdot\|$ -envelopes are also convex, therefore the inclusion  $\mathcal{E}^{k\|\cdot\|} \subset \mathcal{E}^{-k\|\cdot\|}$  is strict. This ensures that the inclusion in (c) of the above theorem generally fails to be an equality.

As regards the function  $\varphi = -k\|\cdot\|^\alpha$  it is also worth mentioning that, for  $\eta(x, y) := \|x - y\|^\alpha$  with  $\alpha > 0$  (even with more general coupling functions) and taking

$$\Phi := \{r - \sigma\eta(\cdot, y) : r \in \mathbb{R}, \sigma > 0, y \in X\},$$

a lower semicontinuous function on the normed space  $X$  is shown in [7, Theorem 4.2] to be  $\Phi$ -convex (i.e. a pointwise supremum of functions in  $\Phi$ ), whenever it is bounded from below by a function in  $\Phi$ . The latter property with  $\alpha = 2$  was previously proved in [29, Theorem 2]. The function  $(x, y) \mapsto -k\|x - y\|^\alpha$  is also used as a particular important example of coupling functions arising in the framework of generalized conjugacy in many papers, see, for example, [24, p. 204].

*Remark 6.1.* Given a non-increasing convex function  $\theta : \mathbb{R}_+ \rightarrow \mathbb{R}$  such that  $\theta(0) = 0$ , one can easily check that the function  $\theta(\|\cdot\|)$  is superadditive. Hence, the previous example can be generalized by taking  $\varphi = \theta(\|\cdot\|)$ .

*Example 6.2.* Let  $X$  be a vector space. Let  $\Lambda \subset X$  be a set containing the origin and such that  $\Lambda + \Lambda \subset \Lambda$ . The function  $\delta_\Lambda$  is clearly subadditive. This implies that the function  $\varphi = -\delta_\Lambda$  is superadditive. By Theorem 6.1(a) it follows that

$$\begin{aligned} f \in \mathcal{E}^{-\delta_\Lambda} &\iff f(x) \geq f(y) \dot{+} (-\delta_\Lambda)(x - y) \quad \text{for all } x, y \in X \\ &\iff f(x) \geq f(y) \quad \text{if } x - y \in \Lambda \\ &\iff f \text{ is } \Lambda\text{-non-decreasing.} \end{aligned}$$

This and Theorem 6.1(b) entail that  $f \nabla \delta_{-\Lambda}$  (respectively  $f \Delta (-\delta_\Lambda)$ ) is the greatest (respectively lowest)  $\Lambda$ -non-decreasing function that is majorized (respectively minorized) by  $f$ . Further, Theorem 6.1(c) says that  $\mathcal{E}^{\delta_\Lambda} \subset \mathcal{E}^{(-\delta_\Lambda)-} = \mathcal{E}^{-\delta_{-\Lambda}}$ , hence the functions of  $\mathcal{E}^{\delta_\Lambda}$  are  $\Lambda$ -non-increasing. In fact, this can be recovered directly by using the characterization of  $\mathcal{E}^{\delta_\Lambda}$  given by Proposition 4.4(ii).

§7. Case  $\varphi \in \Gamma(X)$ .

7.1. Expressions of  $\varphi$ -envelopes as Legendre–Fenchel conjugates. Let us start with the following elementary lemma.

LEMMA 7.1. Let  $X$  be a locally convex space. For every function  $f : X \rightarrow \overline{\mathbb{R}}$ , we have  $(f^*)_- = (f_-)^*$ .

*Proof.* It suffices to use the definition of the Legendre–Fenchel conjugate. For every  $\xi^* \in X^*$ , we have

$$\begin{aligned} (f^*)_-(\xi^*) &= (f^*)(-\xi^*) = \sup_{x \in X} \{ \langle -\xi^*, x \rangle - f(x) \} \\ &= \sup_{y \in X} \{ \langle \xi^*, y \rangle - f(-y) \} \\ &= \sup_{y \in X} \{ \langle \xi^*, y \rangle - f_-(y) \} = (f_-)^*(\xi^*). \quad \square \end{aligned}$$

THEOREM 7.1. Let  $X$  be a locally convex space. Let us assume that  $\varphi \in \Gamma(X)$  and let  $\psi : X^* \rightarrow \overline{\mathbb{R}}$  be such that  $\psi^* = \varphi$ . Then we have for every function  $f : X \rightarrow \overline{\mathbb{R}}$ ,

$$f^\varphi = (\psi \dot{-} (f_-)^*)^*. \tag{43}$$

Moreover, the following equivalences hold:

$$\begin{aligned} g \in \mathcal{E}^\varphi &\iff g = (\psi \dot{-} h)^* \quad \text{for some } h \in \Gamma(X^*) \\ &\iff g = (\psi \dot{-} (\psi \dot{-} g^*)^{**})^*. \end{aligned}$$

*Proof.* For every  $x \in X$ ,

$$\begin{aligned} f^\varphi(x) &= \sup_{y \in X} \{ \varphi(x - y) \dot{-} f(y) \} \\ &= \sup_{y \in X} \left\{ \sup_{\xi^* \in X^*} \{ \langle \xi^*, x - y \rangle - \psi(\xi^*) \} \dot{-} f(y) \right\} \quad \text{since } \varphi = \psi^* \\ &= \sup_{y \in X} \sup_{\xi^* \in X^*} \{ \langle \xi^*, x - y \rangle - \psi(\xi^*) \dot{-} f(y) \} \\ &= \sup_{\xi^* \in X^*} \sup_{y \in X} \{ \langle \xi^*, x - y \rangle - \psi(\xi^*) \dot{-} f(y) \} \\ &= \sup_{\xi^* \in X^*} \left\{ \sup_{y \in X} \{ \langle \xi^*, -y \rangle - f(y) \} \dot{-} \psi(\xi^*) + \langle \xi^*, x \rangle \right\} \\ &= \sup_{\xi^* \in X^*} \{ f^*(-\xi^*) \dot{-} \psi(\xi^*) + \langle \xi^*, x \rangle \} \\ &= (\psi \dot{-} (f^*)_-)^*(x) \\ &= (\psi \dot{-} (f_-)^*)^*(x) \quad \text{in view of Lemma 7.1.} \end{aligned}$$

For the first equivalence, recall that  $g \in \mathcal{E}^\varphi$  if and only if there exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $g = f^\varphi$ . Then use the equality  $f^\varphi = (\psi \dot{-} (f_-)^*)^*$  and the fact that the range of the Legendre–Fenchel transform is equal to  $\Gamma(X^*)$ , see, for example, [22].

For the second equivalence, observe that

$$\begin{aligned} g \in \mathcal{E}^\varphi &\iff g = (g^{\varphi_-})^\varphi \\ &\iff g = [(\psi_- \dot{-} (g_-)^*)^*]^\varphi \quad \text{from formula (43)} \\ &\iff g = [((\psi \dot{-} g^*)^*)_-]^\varphi \quad \text{by Lemma 7.1} \\ &\iff g = (\psi \dot{-} (\psi \dot{-} g^*)^{**})^* \quad \text{from formula (43) again.} \quad \square \end{aligned}$$

*Remark 7.1.* Since  $\varphi \in \Gamma(X)$  by assumption, we have  $\varphi^{**} = \varphi$ , hence we can take  $\psi = \varphi^*$  in the statement of Theorem 7.1.

*Remark 7.2.* Formula (43) can be recovered partially by using a formula on the conjugate of the difference of functions. Recall that for  $\psi : X \rightarrow \mathbb{R} \cup \{+\infty\}$  and  $h \in \Gamma_0(X)$ ,

$$\begin{aligned} \text{for all } x^* \in X^*, \quad (\psi \dot{-} h)^*(x^*) &= \sup_{y^* \in \text{dom } h^*} \{\psi^*(x^* + y^*) - h^*(y^*)\} \\ &= (\psi^* \ominus h^*)(x^*). \end{aligned} \tag{44}$$

This formula is due to Hiriart-Urruty [11]. It was established first by Pshenichnyi [27], assuming that both  $\psi$  and  $h$  are finite-valued convex functions. Now let  $\varphi \in \Gamma_0(X)$  and  $f \in \Gamma_0(X)$ . By reversing the roles of  $X$  and  $X^*$  and by using equality (44) with  $h = (f_-)^*$  and  $\psi : X^* \rightarrow \mathbb{R} \cup \{+\infty\}$  such that  $\psi^* = \varphi$ , we find

$$\begin{aligned} (\psi \dot{-} (f_-)^*)^* &= \varphi \ominus (f_-)^{**} \\ &= \varphi \ominus f_- = f^\varphi. \end{aligned}$$

Hence, we recover formula (43) in the case where both functions  $\varphi$  and  $f$  are in  $\Gamma_0(X)$ .

The next corollary says, in particular, that the  $\varphi$ -envelope of a function coincides with the  $\varphi$ -envelope of its lower semicontinuous convex hull whenever  $\varphi \in \Gamma(X)$ .

**COROLLARY 7.1.** *Let  $X$  be a locally convex space and  $\varphi \in \Gamma(X)$ . Then we have for every function  $f : X \rightarrow \overline{\mathbb{R}}$  and every function  $g : X \rightarrow \overline{\mathbb{R}}$  satisfying  $\overline{\text{co}} f \leq g \leq f$ ,*

$$f^\varphi = (\overline{\text{co}} f)^\varphi = g^\varphi.$$

*Proof.* For the first equality, it suffices to use Theorem 7.1 and the fact that the functions  $f$  and  $\overline{\text{co}} f$  have the same Legendre–Fenchel conjugate. On the other hand, since  $\overline{\text{co}} f \leq g \leq f$ , we see that  $f^\varphi \leq g^\varphi \leq (\overline{\text{co}} f)^\varphi$ . Recalling that  $f^\varphi = (\overline{\text{co}} f)^\varphi$ , the second equality immediately follows. □

For every set  $D \subset X^*$ , we define as in §3 the classes  $\Sigma_D$  and  $\Sigma_D^*$  by

$$\Sigma_D = \{f : X^* \rightarrow \overline{\mathbb{R}}, \text{dom } f \subset D\} \quad \text{and} \quad \Sigma_D^* = \{f^*, f \in \Sigma_D\}.$$

In the same vein, let us define the classes  $\widehat{\Sigma}_D$  and  $\widehat{\Sigma}_D^*$  by

$$\widehat{\Sigma}_D = \{f : X^* \rightarrow \overline{\mathbb{R}}, \text{dom } f = D\} \quad \text{and} \quad \widehat{\Sigma}_D^* = \{f^*, f \in \widehat{\Sigma}_D\}.$$

The following proposition allows us to characterize the classes  $\widehat{\Sigma}_D^*$  and  $\Sigma_D^*$ .

**PROPOSITION 7.1.** *Let  $X$  be a locally convex space and let  $D \subset X^*$  be such that  $D = \{a_i^*, i \in I\}$  for some set  $I$ . Then for every function  $f : X \rightarrow \overline{\mathbb{R}}$ , we have  $f \in \widehat{\Sigma}_D^*$  (respectively  $\Sigma_D^*$ ) if and only if there exists a family  $(\alpha_i)_{i \in I} \subset \mathbb{R} \cup \{+\infty\}$  (respectively  $\overline{\mathbb{R}}$ ) such that  $f = \sup_{i \in I} \langle a_i^*, \cdot \rangle + \alpha_i$ .*

*Proof.* Assume that  $f \in \widehat{\Sigma}_D^*$  (respectively  $\Sigma_D^*$ ). By definition, there exists  $g : X^* \rightarrow \overline{\mathbb{R}}$  such that  $f = g^*$  and  $\text{dom } g = D$  (respectively  $\text{dom } g \subset D$ ). Hence, we have

$$f = \sup_{x^* \in D} \langle x^*, \cdot \rangle - g(x^*) = \sup_{i \in I} \langle a_i^*, \cdot \rangle - g(a_i^*).$$

By setting  $\alpha_i = -g(a_i^*)$  for every  $i \in I$ , we obtain  $f = \sup_{i \in I} \langle a_i^*, \cdot \rangle + \alpha_i$  with  $\alpha_i \in \mathbb{R} \cup \{+\infty\}$  (respectively  $\overline{\mathbb{R}}$ ).

Conversely, assume that there exists  $(\alpha_i)_{i \in I} \subset \mathbb{R} \cup \{+\infty\}$  (respectively  $\overline{\mathbb{R}}$ ) such that  $f = \sup_{i \in I} \langle a_i^*, \cdot \rangle + \alpha_i$ . Then we have

$$\begin{aligned} f &= \sup_{x^* \in D} \left[ \sup_{\{i \in I, a_i^* = x^*\}} \langle a_i^*, \cdot \rangle + \alpha_i \right] \\ &= \sup_{x^* \in D} \left[ \langle x^*, \cdot \rangle + \sup_{\{i \in I, a_i^* = x^*\}} \alpha_i \right]. \end{aligned}$$

Defining the function  $h : X^* \rightarrow \overline{\mathbb{R}}$  by

$$h(x^*) = \begin{cases} \sup_{\{i \in I, a_i^* = x^*\}} \alpha_i & \text{if } x^* \in D, \\ -\infty & \text{if } x^* \notin D, \end{cases}$$

we obtain

$$\begin{aligned} f &= \sup_{x^* \in D} \langle x^*, \cdot \rangle + h(x^*) \\ &= \sup_{x^* \in X^*} \langle x^*, \cdot \rangle + h(x^*). \end{aligned}$$

We conclude that  $f = (-h)^*$  with  $\text{dom}(-h) = D$  (respectively  $\text{dom}(-h) \subset D$ ), hence  $f \in \widehat{\Sigma}_D^*$  (respectively  $f \in \Sigma_D^*$ ). □

*Example 7.1.* Take  $D = \{a_1^*, \dots, a_n^*\} \subset X^*$  for some  $n \geq 1$ . The previous proposition shows that, for every function  $f : X \rightarrow \overline{\mathbb{R}}$ ,

$$f \in \Sigma_D^* \iff f = \sup_{i \in \{1, \dots, n\}} \langle a_i^*, \cdot \rangle + \alpha_i \quad \text{for some } \alpha_1, \dots, \alpha_n \in \overline{\mathbb{R}}.$$

(45)

On the other hand, if  $f \in \Gamma_0(X)$ , the following equivalence holds true:

$$\text{dom } f^* \subset D \iff \text{dom } f^* \subset \{a_i^*\} \text{ for some } i \in \{1, \dots, n\}$$

because the set  $\text{dom } f^*$  is convex. Since  $f^*$  is proper, this is, in turn, equivalent to  $f^* = \delta_{\{a_i^*\}} - \alpha_i$  for some  $\alpha_i \in \mathbb{R}$ . Taking the conjugate, we find  $f = \langle a_i^*, \cdot \rangle + \alpha_i$ . It ensues that the set  $\{f \in \Gamma_0(X), \text{dom } f^* \subset D\}$  coincides with the set of affine continuous functions with slopes in  $D = \{a_1^*, \dots, a_n^*\}$ . This yields an example for which the inclusion (14) is strict. By applying again Proposition 7.1, we obtain that

$$f \in \Sigma_{\text{co}(D)}^* \iff f = \sup_{x^* \in \text{co}(D)} \langle x^*, \cdot \rangle + \alpha_{x^*}, \tag{46}$$

with  $\alpha_{x^*} \in \overline{\mathbb{R}}$  for every  $x^* \in \text{co}(D)$ . The comparison of (45) and (46) clearly shows that the inclusion  $\Sigma_D^* \subset \Sigma_{\text{co}(D)}^*$  is strict as soon as the set  $D = \{a_1^*, \dots, a_n^*\}$  is not a singleton. This easily implies that the inclusion (15) is strict for such a set  $D$ .

The next result gives several upper bounds (in the sense of inclusion) for the set  $\mathcal{E}^\varphi$ , respectively when  $\varphi \in \Gamma(X)$ ,  $\varphi \in \widehat{\Sigma}_D^*$  and  $\varphi \in \Sigma_D^*$ .

**COROLLARY 7.2.** *Let  $X$  be a locally convex space and let  $\varphi \in \Gamma(X)$ .*

(i) *The following inclusions hold true:*

$$\mathcal{E}^\varphi \subset \bigcap_{\{\psi, \varphi = \psi^*\}} (\widehat{\Sigma}_{\text{dom } \psi}^* \cup \{-\omega_X\}) \subset \bigcap_{\{\psi, \varphi = \psi^*\}} \Sigma_{\text{dom } \psi}^*. \tag{47}$$

(ii) *For every subset  $D \subset X^*$ , we have*

$$\begin{aligned} \varphi \in \widehat{\Sigma}_D^* &\iff \mathcal{E}^\varphi \subset \widehat{\Sigma}_D^* \cup \{-\omega_X\} \text{ if } \varphi \neq -\omega_X, \\ \varphi \in \Sigma_D^* &\iff \mathcal{E}^\varphi \subset \Sigma_D^*. \end{aligned}$$

(iii) *Assume that there exist families  $(a_i^*)_{i \in I} \subset X^*$  and  $(\alpha_i)_{i \in I} \subset \mathbb{R} \cup \{+\infty\}$  (respectively  $\overline{\mathbb{R}}$ ) such that*

$$\varphi = \sup_{i \in I} \langle a_i^*, \cdot \rangle + \alpha_i.$$

*Then for every  $g \in \mathcal{E}^\varphi \setminus \{-\omega_X\}$  (respectively  $g \in \mathcal{E}^\varphi$ ), there exists  $(\beta_i)_{i \in I} \subset \mathbb{R} \cup \{+\infty\}$  (respectively  $\overline{\mathbb{R}}$ ) such that*

$$g = \sup_{i \in I} \langle a_i^*, \cdot \rangle + \beta_i.$$

*In particular, if the set  $I$  is finite, every  $\varphi$ -envelope is polyhedral.*

*Proof.* (i) Let  $\psi : X^* \rightarrow \overline{\mathbb{R}}$  be such that  $\varphi = \psi^*$ . Assuming that  $g \in \mathcal{E}^\varphi$ , Theorem 7.1 shows that  $g = (\psi \dot{-} h)^*$  for some  $h \in \Gamma(X^*)$ . If  $h = -\omega_{X^*}$ , we have  $\psi \dot{-} h = \omega_{X^*}$  and therefore  $g = -\omega_X$ . If  $h \neq -\omega_{X^*}$ , we see that  $\text{dom}(\psi \dot{-} h) = \text{dom } \psi$ , hence  $g \in \widehat{\Sigma}_{\text{dom } \psi}^*$ . We deduce the inclusion

$\mathcal{E}^\varphi \subset \widehat{\Sigma}_{\text{dom } \psi}^* \cup \{-\omega_X\}$ . Since this is true for every function  $\psi : X^* \rightarrow \overline{\mathbb{R}}$  such that  $\varphi = \psi^*$ , the first inclusion of (47) follows. For the second inclusion, it suffices to note that  $\widehat{\Sigma}_{\text{dom } \psi}^* \cup \{-\omega_X\} \subset \Sigma_{\text{dom } \psi}^*$ .

(ii) Let us fix  $D \subset X^*$  and assume that  $\varphi \in \widehat{\Sigma}_D^*$ . Then there exists  $\psi : X^* \rightarrow \overline{\mathbb{R}}$  such that  $\varphi = \psi^*$  and  $\text{dom } \psi = D$ . We deduce from the first inclusion of (47) that

$$\mathcal{E}^\varphi \subset \widehat{\Sigma}_{\text{dom } \psi}^* \cup \{-\omega_X\} = \widehat{\Sigma}_D^* \cup \{-\omega_X\}.$$

Conversely, if  $\mathcal{E}^\varphi \subset \widehat{\Sigma}_D^* \cup \{-\omega_X\}$  and if  $\varphi \neq -\omega_X$ , then we obtain  $\varphi \in \widehat{\Sigma}_D^*$  according to the inclusion  $\varphi \in \mathcal{E}^\varphi$ . The proof of the second equivalence is analogous and left to the reader.

(iii) Let  $(a_i^*)_{i \in I} \subset X^*$  and  $(\alpha_i)_{i \in I} \subset \mathbb{R} \cup \{+\infty\}$  (respectively  $\overline{\mathbb{R}}$ ) be such that  $\varphi = \sup_{i \in I} \langle a_i^*, \cdot \rangle + \alpha_i$ . Let us set  $D = \{a_i^*, i \in I\}$ . Proposition 7.1 shows that  $\varphi \in \widehat{\Sigma}_D^*$  (respectively  $\Sigma_D^*$ ). If  $g \in \mathcal{E}^\varphi \setminus \{-\omega_X\}$  (respectively  $g \in \mathcal{E}^\varphi$ ), we deduce from (ii) that  $g \in \widehat{\Sigma}_D^*$  (respectively  $\Sigma_D^*$ ). By applying Proposition 7.1 again, we derive the existence of  $(\beta_i)_{i \in I} \subset \mathbb{R} \cup \{+\infty\}$  (respectively  $\overline{\mathbb{R}}$ ) such that  $g = \sup_{i \in I} \langle a_i^*, \cdot \rangle + \beta_i$ . Finally, if the set  $I$  is finite and if  $g$  is a  $\varphi$ -envelope, then either  $g = \pm\omega_X$  or the function  $g$  is the supremum of a finite collection of continuous affine functions. We then conclude that  $g$  is polyhedral.  $\square$

By applying the second equivalence of Corollary 7.2(ii) with  $D = X^*$ , we obtain that  $\varphi \in \Gamma(X)$  if and only if  $\mathcal{E}^\varphi \subset \Gamma(X)$ . Corollary 7.3 below shows that in this case the set  $\mathcal{E}^\varphi$  is strictly included in  $\Gamma(X)$ . Note that for  $\varphi \in \Gamma_0(X)$  satisfying a suitable condition (named generating condition), the functions of the class  $\mathcal{E}^\varphi$  have been studied in [25] under the terminology of  $\varphi$ -strongly convex functions.

Following Theorem 7.1 and Remark 7.1, we have  $g \in \mathcal{E}^\varphi$  if and only if  $g = (\varphi^* \dot{-} h)^*$  for some  $h \in \Gamma(X^*)$ . Let us now have a look at the class of the functions equal to  $(\varphi^* \dot{-} h)^*$  for some  $h : X^* \rightarrow \mathbb{R} \cup \{+\infty\}$  not necessarily in  $\Gamma(X^*)$ .

**PROPOSITION 7.2.** *Let  $X$  be a locally convex space. Assume that  $\varphi \in \Gamma_0(X)$  and  $g \in \Gamma_0(X)$ .*

- (i) *If  $g = (\varphi^* \dot{-} h)^*$  for some  $h : X^* \rightarrow \mathbb{R} \cup \{+\infty\}$ , then we have  $g^\infty = \varphi^\infty$ , which is equivalent to  $\text{cl}^{w^*}(\text{dom } g^*) = \text{cl}^{w^*}(\text{dom } \varphi^*)$ .*
- (ii) *If  $\text{dom } g^* = \text{dom } \varphi^*$ , then  $g = (\varphi^* \dot{-} h)^*$  for  $h : X^* \rightarrow \mathbb{R} \cup \{+\infty\}$  given by  $h = \varphi^* \dot{-} g^*$ .*

*Proof.* (i) Assume that  $g = (\varphi^* \dot{-} h)^*$  for some  $h : X^* \rightarrow \mathbb{R} \cup \{+\infty\}$ . By definition of the Legendre–Fenchel transform, we obtain

$$\begin{aligned} g &= \sup_{\xi^* \in X^*} \{ \langle \xi^*, \cdot \rangle + h(\xi^*) - \varphi^*(\xi^*) \} \\ &= \sup_{\xi^* \in \text{dom } \varphi^*} \{ \langle \xi^*, \cdot \rangle + h(\xi^*) - \varphi^*(\xi^*) \}. \end{aligned} \tag{48}$$

Observe that the function  $h$  cannot take the value  $+\infty$  on  $\text{dom } \varphi^*$  (otherwise we would have  $g = \omega_X$ ). Therefore, the values  $-\varphi^*(\xi^*)$  and  $h(\xi^*)$  are finite for every  $\xi^* \in \text{dom } \varphi^*$ . By taking the recession function of each member of (48), we obtain

$$g^\infty = \sup_{\xi^* \in \text{dom } \varphi^*} [(\xi^*, \cdot) + h(\xi^*) - \varphi^*(\xi^*)]^\infty.$$

The recession function of the affine map  $(\xi^*, \cdot) + h(\xi^*) - \varphi^*(\xi^*)$  is equal to  $(\xi^*, \cdot)$ , thus implying that  $g^\infty = \sup_{\xi^* \in \text{dom } \varphi^*} (\xi^*, \cdot) = \sigma_{\text{dom } \varphi^*}$ . Recalling that  $\sigma_{\text{dom } \varphi^*} = \varphi^\infty$ , we deduce that  $g^\infty = \varphi^\infty$ , which is in turn equivalent to the equality  $\text{cl}^{w*}(\text{dom } g^*) = \text{cl}^{w*}(\text{dom } \varphi^*)$ , see [22].

(ii) Assume that  $\text{dom } g^* = \text{dom } \varphi^*$ . It is easy to check that for every  $x^* \in X^*$ ,

$$(\varphi^* \dot{-} (\varphi^* \dot{-} g^*))(x^*) = \begin{cases} g^*(x^*) & \text{if } x^* \in \text{dom } g^*, \\ +\infty & \text{if } x^* \notin \text{dom } g^*. \end{cases}$$

It ensues that  $\varphi^* \dot{-} (\varphi^* \dot{-} g^*) = g^*$ . Since  $g \in \Gamma_0(X)$  by assumption, we have  $g = g^{**}$ , hence  $g = (\varphi^* \dot{-} (\varphi^* \dot{-} g^*))^*$ . The function  $h = \varphi^* \dot{-} g^*$  takes its values in  $\mathbb{R} \cup \{+\infty\}$  because  $\text{dom } g^* = \text{dom } \varphi^*$ . □

Combining Theorem 7.1 and Proposition 7.2, we derive a necessary (respectively sufficient) condition for a function  $g \in \Gamma_0(X)$  to be a  $\varphi$ -envelope.

**COROLLARY 7.3.** *Let  $X$  be a locally convex space. Assume that  $\varphi \in \Gamma_0(X)$  and  $g \in \Gamma_0(X)$ .*

- (i) *If  $g \in \mathcal{E}^\varphi$  then  $g^\infty = \varphi^\infty$ .*
- (ii) *If  $\text{dom } g^* = \text{dom } \varphi^*$  and  $\varphi^* \dot{-} g^* \in \Gamma_0(X^*)$ , then  $g \in \mathcal{E}^\varphi$ .*

*Proof.* (i) If  $g \in \mathcal{E}^\varphi$ , we deduce from Theorem 7.1 that  $g = (\varphi^* \dot{-} h)^*$  for some  $h \in \Gamma(X^*)$ . Since  $g \in \Gamma_0(X)$  by assumption, we have  $h \neq -\omega_{X^*}$ , hence the function  $h$  does not take the value  $-\infty$ . Proposition 7.2(i) then implies that  $g^\infty = \varphi^\infty$ .

(ii) If  $\text{dom } g^* = \text{dom } \varphi^*$ , Proposition 7.2(ii) shows that  $g = (\varphi^* \dot{-} h)^*$  with  $h = \varphi^* \dot{-} g^*$ . Since  $h \in \Gamma_0(X^*)$  by assumption, we conclude by Theorem 7.1 that  $g \in \mathcal{E}^\varphi$ . □

**7.2. Klee envelopes.** Let  $(X, \|\cdot\|)$  be a normed space and let  $f : X \rightarrow \overline{\mathbb{R}}$  be an extended real-valued function. For any reals  $\lambda > 0$  and  $p \geq 1$ , we define the Klee envelope of  $f$  with index  $\lambda$  and power  $p$  as

$$\kappa_{\lambda,p} f(x) = \sup_{y \in X} \left( \frac{1}{p\lambda} \|x - y\|^p - f(y) \right).$$



In other words, we have  $\kappa_{\lambda,p}f = f^\varphi$  with the function  $\varphi : X \rightarrow \mathbb{R}$  defined by  $\varphi(x) = (1/p\lambda)\|x\|^p$ . Applying Theorem 7.1 with  $\varphi = (1/p\lambda)\|\cdot\|^p$  and denoting by  $\|\cdot\|_{X^*}$  the dual norm on  $X^*$  we obtain the following result.

**COROLLARY 7.4.** *Let  $(X, \|\cdot\|)$  be a normed space. For any  $\lambda > 0, p > 1$  and for every function  $f : X \rightarrow \mathbb{R}$ , we have*

$$\kappa_{\lambda,p}f = \left( \frac{\lambda^{q-1}}{q} \|\cdot\|_{X^*}^q - (f-\cdot)^* \right)^*, \tag{49}$$

where  $q > 1$  is the conjugate exponent of  $p$ . Moreover, the following assertions are equivalent:

- (i)  $g$  is a Klee envelope with index  $\lambda$  and power  $p$ ;
- (ii)  $g = ((\lambda^{q-1}/q)\|\cdot\|_{X^*}^q - h)^*$  for some  $h \in \Gamma(X^*)$ ;
- (iii)  $g = ((\lambda^{q-1}/q)\|\cdot\|_{X^*}^q - ((\lambda^{q-1}/q)\|\cdot\|_{X^*}^q - g^*)^{**})^*$ .

These assertions are satisfied whenever the following stronger condition is fulfilled:

- (iv)  $g \in \Gamma(X)$  and  $(\lambda^{q-1}/q)\|\cdot\|_{X^*}^q - g^* \in \Gamma(X^*)$ .

*Proof.* It suffices to apply Theorem 7.1 with  $\varphi = (1/p\lambda)\|\cdot\|^p$  and  $\psi = \varphi^* = (\lambda^{q-1}/q)\|\cdot\|_{X^*}^q$ . Let us now establish the implication (iv)  $\implies$  (ii). Assume that  $g \in \Gamma(X)$  and that  $(\lambda^{q-1}/q)\|\cdot\|_{X^*}^q - g^* \in \Gamma(X^*)$ . The function  $g^*$  can be written as  $g^* = (\lambda^{q-1}/q)\|\cdot\|_{X^*}^q - h$  for some  $h \in \Gamma(X^*)$ . Since  $g \in \Gamma(X)$  by assumption, we have  $g = g^{**}$ . Hence, we deduce that  $g = ((\lambda^{q-1}/q)\|\cdot\|_{X^*}^q - h)^*$  and assertion (ii) is proved.  $\square$

**COROLLARY 7.5.** *Let  $(X, \|\cdot\|)$  be a normed space. For every  $p > 1$  and every  $C \subset X$ , the farthest distance function  $\Delta_C = \sup_{y \in C} \|\cdot - y\|$  satisfies*

$$\frac{1}{p} \Delta_C^p = \left( \frac{1}{q} \|\cdot\|_{X^*}^q - \sigma_{-C} \right)^*.$$

*Proof.* Observe that

$$\kappa_{1,p} \delta_C = \sup_{y \in X} \left\{ \frac{1}{p} \|\cdot - y\|^p - \delta_C(y) \right\} = \sup_{y \in C} \frac{1}{p} \|\cdot - y\|^p = \frac{1}{p} \Delta_C^p.$$

It suffices then to apply formula (49) of Corollary 7.4 with  $f = \delta_C$  and  $\lambda = 1$ .  $\square$

Additional properties of the Klee envelopes can be obtained in the case when  $(X, \|\cdot\|)$  is a Hilbert space and  $p = 2$ .

**THEOREM 7.2.** *Assume that  $X$  is a Hilbert space endowed with the scalar product  $\langle \cdot, \cdot \rangle$  and the corresponding norm  $\|\cdot\|$ .*

(a) For every  $\lambda > 0$  and every function  $f : X \rightarrow \overline{\mathbb{R}}$ , we have

$$\kappa_{\lambda,2}f = \left( \frac{\lambda}{2} \|\cdot\|^2 - (f_-)^* \right)^* \tag{50}$$

$$= \left( f - \frac{1}{2\lambda} \|\cdot\|^2 \right)^* \left( -\frac{\cdot}{\lambda} \right) + \frac{1}{2\lambda} \|\cdot\|^2; \tag{51}$$

$$\kappa_{\lambda,2}(\kappa_{\lambda,2}f) = \left( f - \frac{1}{2\lambda} \|\cdot\|^2 \right)^{**} + \frac{1}{2\lambda} \|\cdot\|^2. \tag{52}$$

(b) For  $\lambda > 0$  and  $f : X \rightarrow \overline{\mathbb{R}}$  the following assertions are equivalent:

- (i)  $f$  is a Klee envelope with index  $\lambda$  and power 2;
- (ii)  $f = ((\lambda/2)\|\cdot\|^2 - h)^*$  for some  $h \in \Gamma(X)$ ;
- (iii)  $f = ((\lambda/2)\|\cdot\|^2 - ((\lambda/2)\|\cdot\|^2 - f^*)^{**})^*$ ;
- (iv)  $f - (1/2\lambda)\|\cdot\|^2 \in \Gamma(X)$ ;
- (v)  $f \in \Gamma(X)$  and  $(\lambda/2)\|\cdot\|^2 - f^* \in \Gamma(X)$ .

*Proof.* (a) For the equality (50), it suffices to apply Corollary 7.4 with  $p = 2$ . For the equality (51), observe that for every  $x \in X$ ,

$$\begin{aligned} \kappa_{\lambda,2}f(x) &= \sup_{y \in X} \left\{ \frac{1}{2\lambda} \|x - y\|^2 - f(y) \right\} \\ &= \sup_{y \in X} \left\{ \frac{1}{2\lambda} \|x\|^2 + \frac{1}{2\lambda} \|y\|^2 - \frac{1}{\lambda} \langle x, y \rangle - f(y) \right\} \\ &= \left( f - \frac{1}{2\lambda} \|\cdot\|^2 \right)^* (-x/\lambda) + \frac{1}{2\lambda} \|x\|^2. \end{aligned}$$

By iterating we deduce that

$$\begin{aligned} \kappa_{\lambda,2}(\kappa_{\lambda,2}f) &= \left( \kappa_{\lambda,2}f - \frac{1}{2\lambda} \|\cdot\|^2 \right)^* \left( -\frac{\cdot}{\lambda} \right) + \frac{1}{2\lambda} \|\cdot\|^2 \\ &= \left[ \left( f - \frac{1}{2\lambda} \|\cdot\|^2 \right)^* \left( -\frac{\cdot}{\lambda} \right) \right]^* \left( -\frac{\cdot}{\lambda} \right) + \frac{1}{2\lambda} \|\cdot\|^2 \\ &= \left( f - \frac{1}{2\lambda} \|\cdot\|^2 \right)^{**} + \frac{1}{2\lambda} \|\cdot\|^2, \end{aligned}$$

which proves the equality (52).

(b) We now show that assertions (i) to (v) are equivalent. The equivalences (i)  $\iff$  (ii)  $\iff$  (iii) are consequences of Corollary 7.4 applied with  $p = 2$ . Let us show the equivalence (i)  $\iff$  (iv). Observe that  $f$  is a Klee envelope with index  $\lambda$  and power 2 if and only if  $f \in \mathcal{E}^\varphi$  with  $\varphi = (1/2\lambda)\|\cdot\|^2$ . From the equivalence (7)  $\iff$  (8) and the fact that  $\varphi_- = \varphi$ , this is, in turn, equivalent to

$f = (f^\varphi)^\varphi$ . Since  $(f^\varphi)^\varphi = \kappa_{\lambda,2}(\kappa_{\lambda,2}f)$  and using the equality (52), we infer that

$f$  is a Klee envelope with index  $\lambda$  and power 2

$$\begin{aligned} f - \frac{1}{2\lambda} \|\cdot\|^2 &= \left( f - \frac{1}{2\lambda} \|\cdot\|^2 \right)^{**} \\ f - \frac{1}{2\lambda} \|\cdot\|^2 &\in \Gamma(X). \end{aligned}$$

Hence the equivalence (i)  $\iff$  (iv) is proved. Let us now show that (iv)  $\implies$  (v). If  $f - (1/2\lambda)\|\cdot\|^2 = \pm\omega_X$ , then assertion (v) is trivially satisfied. Hence we can assume that  $f = (1/2\lambda)\|\cdot\|^2 + h$  with  $h \in \Gamma_0(X)$ . This clearly implies that  $f \in \Gamma_0(X)$ . Taking the conjugate, we obtain that  $f^* = (\lambda/2)\|\cdot\|^2 \nabla h^*$  since the classical qualification condition is satisfied. It ensues that for every  $x \in X$ ,

$$\begin{aligned} f^*(x) &= \inf_{y \in X} \left\{ \frac{\lambda}{2} \|x - y\|^2 + h^*(y) \right\} \\ &= \frac{\lambda}{2} \|x\|^2 + \inf_{y \in X} \left\{ -\lambda \langle x, y \rangle + \frac{\lambda}{2} \|y\|^2 + h^*(y) \right\}. \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{\lambda}{2} \|x\|^2 - f^*(x) &= \sup_{y \in X} \left\{ \lambda \langle x, y \rangle - \frac{\lambda}{2} \|y\|^2 - h^*(y) \right\} \\ &= \left( h^* + \frac{\lambda}{2} \|\cdot\|^2 \right)^* (\lambda x). \end{aligned}$$

This clearly implies that  $(\lambda/2)\|\cdot\|^2 - f^* \in \Gamma_0(X)$  and (v) is proved. Let us finally observe that the implication (v)  $\implies$  (ii) has been established in Corollary 7.4. As a conclusion, we have shown the equivalences (i)  $\iff$  (ii)  $\iff$  (iii)  $\iff$  (iv) along with the implications (iv)  $\implies$  (v)  $\implies$  (ii), which clearly establishes that all assertions (i) to (v) are equivalent.  $\square$

The equalities (51) and (52) have been previously established by Wang [39] respectively in Proposition 4.5 and at the end of the proof of Proposition 4.13. As noted in [39, Proposition 4.13] those equalities directly yield, for  $f$  proper and lower semicontinuous, that  $\kappa_{\lambda,2}(\kappa_{\lambda,2}f) = f$  if and only if  $f - (1/2\lambda)\|\cdot\|^2$  is convex.

Taking  $f$  as the indicator function of a set  $C$  gives the following corollary.

**COROLLARY 7.6.** *Assume that  $X$  is a Hilbert space. For every  $C \subset X$ , the farthest distance function  $\Delta_C$  satisfies*

$$\frac{1}{2} \Delta_C^2 = \left( \frac{1}{2} \|\cdot\|^2 - \sigma_{-C} \right)^* = \left( \delta_{-C} - \frac{1}{2} \|\cdot\|^2 \right)^* + \frac{1}{2} \|\cdot\|^2.$$

*Proof.* It suffices to apply formulas (50)–(51) of Theorem 7.2 with  $f = \delta_C$  and  $\lambda = 1$ . □

7.3. *Case of a positively homogeneous function  $\varphi$ .* In this subsection, we assume that  $X$  is a locally convex space and that the function  $\varphi \in \Gamma_0(X)$  is positively homogeneous, i.e.  $\varphi = \sigma_D$  for a non-empty set  $D \subset X^*$ . By applying Theorem 7.1 with  $\psi = \delta_D$ , we immediately obtain the following result.

**COROLLARY 7.7.** *Let  $X$  be a locally convex space. Take  $\varphi = \sigma_D$  for a non-empty set  $D \subset X^*$ . Then we have, for every function  $f : X \rightarrow \overline{\mathbb{R}}$ ,*

$$f^\varphi = (\delta_D \dot{-} (f_-)^*)^* = \sup_{\xi^* \in D} \{ \langle \xi^*, \cdot \rangle + f^*(-\xi^*) \}.$$

Moreover,

$$\begin{aligned} g \in \mathcal{E}^\varphi &\iff g = (\delta_D \dot{-} h)^* = \sup_{\xi^* \in D} \{ \langle \xi^*, \cdot \rangle + h(\xi^*) \} \text{ for some } h \in \Gamma(X^*) \\ &\iff g = (\delta_D \dot{-} (\delta_D \dot{-} g^*)^{**})^*. \end{aligned}$$

Let us now particularize to the case of a normed space  $(X, \|\cdot\|)$  and take  $\varphi = \|\cdot\|$ .

**COROLLARY 7.8.** *Let  $(X, \|\cdot\|)$  be a normed space. For every function  $f : X \rightarrow \overline{\mathbb{R}}$ , we have*

$$\begin{aligned} \kappa_{1,1}f &= (\delta_{\mathbb{B}_{X^*}} \dot{-} (f_-)^*)^* = \sup_{\xi^* \in \mathbb{B}_{X^*}} \{ \langle \xi^*, \cdot \rangle + f^*(-\xi^*) \} \\ &= (\delta_{\mathbb{S}_{X^*}} \dot{-} (f_-)^*)^* = \sup_{\xi^* \in \mathbb{S}_{X^*}} \{ \langle \xi^*, \cdot \rangle + f^*(-\xi^*) \}. \end{aligned}$$

Moreover,

*$g$  is a Klee envelope with index 1 and power 1*

$$\begin{aligned} &\Updownarrow \\ g &= (\delta_{\mathbb{B}_{X^*}} \dot{-} h)^* = \sup_{\xi^* \in \mathbb{B}_{X^*}} \{ \langle \xi^*, \cdot \rangle + h(\xi^*) \} \text{ for some } h \in \Gamma(X^*) \\ &\Updownarrow \\ g &= (\delta_{\mathbb{B}_{X^*}} \dot{-} (\delta_{\mathbb{B}_{X^*}} \dot{-} g^*)^{**})^* \\ &\Updownarrow \\ g &= (\delta_{\mathbb{S}_{X^*}} \dot{-} h)^* = \sup_{\xi^* \in \mathbb{S}_{X^*}} \{ \langle \xi^*, \cdot \rangle + h(\xi^*) \} \text{ for some } h \in \Gamma(X^*) \\ &\Updownarrow \\ g &= (\delta_{\mathbb{S}_{X^*}} \dot{-} (\delta_{\mathbb{S}_{X^*}} \dot{-} g^*)^{**})^*. \end{aligned}$$

*Proof.* For the equalities  $\kappa_{1,1}f = (\delta_{\mathbb{B}_{X^*}} \dot{-} (f_-)^*)^*$  and  $\kappa_{1,1}f = (\delta_{\mathbb{S}_{X^*}} \dot{-} (f_-)^*)^*$ , use Corollary 7.7 respectively with  $D = \mathbb{B}_{X^*}$  and  $D = \mathbb{S}_{X^*}$ . The characterizations of Klee envelopes with index 1 and power 1 follow immediately.  $\square$

Assuming that  $f = \delta_C$ , we have

$$\kappa_{1,1}\delta_C = \sup_{x \in X} \{ \| \cdot - x \| - \delta_C(x) \} = \sup_{x \in C} \| \cdot - x \| = \Delta_C,$$

where  $\Delta_C$  is the farthest distance function. Taking into account the previous corollary, we then obtain

$$\Delta_C = (\delta_{\mathbb{B}_{X^*}} \dot{-} \sigma_{-C})^* = (\delta_{\mathbb{S}_{X^*}} \dot{-} \sigma_{-C})^*.$$

It is interesting to compare this expression with that of the signed distance  $\text{sgd}(\cdot, C)$  defined by  $\text{sgd}(\cdot, C) := d(\cdot, C) - d(\cdot, X \setminus C)$ , for which it is known that  $\text{sgd}(\cdot, C) = (\delta_{\mathbb{S}_{X^*}} + \sigma_C)^*$ , see [23].

Consider now the case of a finite set  $D = \{a_1^*, \dots, a_n^*\} \subset X^*$  for  $n \geq 1$ . By applying Corollary 7.7, we obtain the following result.

**COROLLARY 7.9.** *Let  $X$  be a locally convex space. Take  $\varphi = \sigma_{\{a_1^*, \dots, a_n^*\}}$  with  $a_1^*, \dots, a_n^* \in X^*$  and  $n \geq 1$ . Then we have, for every function  $f : X \rightarrow \overline{\mathbb{R}}$ ,*

$$f^\varphi = \sup_{i \in \{1, \dots, n\}} \langle a_i^*, \cdot \rangle + f^*(-a_i^*).$$

Moreover,

$$g \in \mathcal{E}^\varphi \iff g = \sup_{i \in \{1, \dots, n\}} \langle a_i^*, \cdot \rangle + h(a_i^*) \text{ for some } h \in \Gamma(X^*).$$

§8. Case  $\varphi \in -\Gamma(X)$ .

8.1. Links between  $\varphi$ -envelopes and Legendre–Fenchel conjugates.

**PROPOSITION 8.1.** *Let  $X$  be a locally convex space and let  $\varphi, g : X \rightarrow \overline{\mathbb{R}}$  be extended real-valued functions.*

- (i) *If  $g \in \mathcal{E}^\varphi$ , then there exists  $h \in \Gamma(X^*)$  such that  $(-g)^* = (-\varphi)^* \dot{+} h$ . If, in addition,  $g \in -\Gamma(X)$ , then  $-g = ((-\varphi)^* \dot{+} h)^*$ .*
- (ii) *Assume that  $X$  is normed. If  $\varphi \in -\Gamma(X)$  and if there exists  $h \in \Gamma(X^*)$  satisfying the equality  $-g = ((-\varphi)^* \dot{+} h)^*$  along with the condition  $0 \in \text{int}(\text{dom } h - \text{dom}(-\varphi)^*)$ , then  $g \in \mathcal{E}^\varphi$ .*

*Proof.* (i) Since  $g \in \mathcal{E}^\varphi$ , there exists  $f : X \rightarrow \overline{\mathbb{R}}$  such that  $g = f^\varphi$ , hence  $-g = (-\varphi) \nabla f$  by (3). Taking the conjugate of each member, we find  $(-g)^* = (-\varphi)^* \dagger f^*$ . Hence, the expected equality holds with  $h = f^* \in \Gamma(X^*)$ . If, in addition,  $g \in -\Gamma(X)$ , we have  $-g = (-g)^{**}$ , hence we deduce from what precedes that  $-g = ((-\varphi)^* \dagger h)^*$ .

(ii) Assume that  $-g = ((-\varphi)^* \dagger h)^*$  for some  $h \in \Gamma(X^*)$ . If  $h = -\omega_{X^*}$  or if  $(-\varphi)^* = -\omega_{X^*}$ , then  $-g = (-\omega_{X^*})^* = \omega_X$  and the inclusion  $g \in \mathcal{E}^\varphi$  trivially holds. Now assume that  $h \neq -\omega_{X^*}$  and  $(-\varphi)^* \neq -\omega_{X^*}$ . Since  $0 \in \text{int}(\text{dom } h - \text{dom}(-\varphi)^*)$ , the functions  $(-\varphi)^*$  and  $h$  are proper and according to the fact that  $X^*$  is a Banach space, we have

$$\begin{aligned} -g &= (-\varphi)^{**} \nabla h^* \\ &= (-\varphi) \nabla h^* \quad \text{because } \varphi \in -\Gamma(X). \end{aligned}$$

We conclude that  $g = \varphi \Delta (-h^*) = (h^*)^\varphi \in \mathcal{E}^\varphi$ . □

**COROLLARY 8.1.** *Let  $X$  be a normed space and let  $\varphi \in -\Gamma_0(X)$  be such that  $\text{dom}(-\varphi)^* = X^*$ . For every  $g \in -\Gamma(X)$ , the following equivalences hold true:*

$$\begin{aligned} g \in \mathcal{E}^\varphi &\iff (-g)^* - (-\varphi)^* \in \Gamma(X^*) \\ &\iff -g = ((-\varphi)^* \dagger h)^* \quad \text{for some } h \in \Gamma(X^*). \end{aligned}$$

*Proof.* Fix  $g \in -\Gamma(X)$ . Since  $\text{dom}(-\varphi)^* = X^*$  and  $-\varphi \in \Gamma_0(X)$ , the function  $(-\varphi)^*$  is finite-valued on  $X^*$ , so the implication

$$g \in \mathcal{E}^\varphi \implies h := (-g)^* - (-\varphi)^* \in \Gamma(X^*)$$

follows from Proposition 8.1(i). Recalling that  $g \in -\Gamma(X)$ , the right-hand inclusion implies in turn that  $-g = ((-\varphi)^* \dagger h)^*$ .

Now assume that  $-g = ((-\varphi)^* \dagger h)^*$  for some  $h \in \Gamma(X^*)$ . If  $\text{dom } h \neq \emptyset$ , the qualification assumption  $0 \in \text{int}(\text{dom } h - \text{dom}(-\varphi)^*)$  is automatically satisfied. We then deduce from Proposition 8.1(ii) that  $g \in \mathcal{E}^\varphi$ . On the other hand, if  $\text{dom } h = \emptyset$ , then we have  $h = \omega_{X^*}$  and hence  $-g = (\omega_{X^*})^* = -\omega_X$ . Then the inclusion  $g \in \mathcal{E}^\varphi$  trivially holds. □

**8.2. Moreau envelopes.** Let  $(X, \|\cdot\|)$  be a normed space and let  $f : X \rightarrow \overline{\mathbb{R}}$  be an extended real-valued function. For  $\lambda > 0$  and  $p \geq 1$ , we define the Moreau envelope of  $f$  with index  $\lambda$  and power  $p$  as

$$e_{\lambda,p} f = \inf_{y \in X} \left( \frac{1}{p\lambda} \|\cdot - y\|^p + f(y) \right) = \frac{1}{p\lambda} \|\cdot\|^p \nabla f.$$

Observe that  $-e_{\lambda,p} f = (-(1/p\lambda)\|\cdot\|^p) \Delta (-f) = f^\varphi$ , with the function  $\varphi : X \rightarrow \overline{\mathbb{R}}$  defined by  $\varphi = -(1/p\lambda)\|\cdot\|^p$ . It ensues that  $g$  is a Moreau envelope with index  $\lambda$  and power  $p$  if and only if  $-g \in \mathcal{E}^\varphi$ , for  $\varphi = -(1/p\lambda)\|\cdot\|^p$ . By applying the results of the previous subsection with  $\varphi = -(1/p\lambda)\|\cdot\|^p$ , we obtain the following statement.

**COROLLARY 8.2.** *Assume that  $(X, \|\cdot\|)$  is a normed space. Let  $\lambda > 0$ ,  $p > 1$  and let  $q$  be the conjugate exponent of  $p$ .*

- (i) *If  $g$  is a Moreau envelope with index  $\lambda$  and power  $p$ , then the function  $g^* - (\lambda^{q-1}/q)\|\cdot\|_{X^*}^q \in \Gamma(X^*)$ .*
- (ii) *If moreover  $g \in \Gamma(X)$ , the following equivalences hold true:*

$$\begin{aligned}
 &g \text{ is a Moreau envelope with index } \lambda \text{ and power } p \\
 &\quad \Updownarrow \\
 &g^* - \frac{\lambda^{q-1}}{q} \|\cdot\|_{X^*}^q \in \Gamma(X^*) \\
 &\quad \Updownarrow \\
 &g = \left( \frac{\lambda^{q-1}}{q} \|\cdot\|_{X^*}^q + h \right)^* \text{ for some } h \in \Gamma(X^*).
 \end{aligned}$$

*Proof.* (i) It suffices to apply Proposition 8.1(i) with  $\varphi = -(1/p\lambda)\|\cdot\|^p$  and to recall that  $((1/p\lambda)\|\cdot\|^p)^* = (\lambda^{q-1}/q)\|\cdot\|_{X^*}^q$ .  
 (ii) The equivalences follow from Corollary 8.1 applied with

$$\varphi = -\frac{1}{p\lambda} \|\cdot\|^p. \quad \square$$

When  $X$  is a Hilbert space, we obtain a more precise characterization of Moreau envelopes with power 2, as shown by the following proposition.

**PROPOSITION 8.2.** *Assume that  $X$  is a Hilbert space endowed with the scalar product  $\langle \cdot, \cdot \rangle$  and the corresponding norm  $\|\cdot\|$ .*

- (a) *For every  $\lambda > 0$  and every function  $f : X \rightarrow \overline{\mathbb{R}}$ , we have*

$$e_{\lambda,2}f = -\left(f + \frac{1}{2\lambda} \|\cdot\|^2\right)^* \left(\frac{\cdot}{\lambda}\right) + \frac{1}{2\lambda} \|\cdot\|^2. \tag{53}$$

*The  $\lambda$ -proximal hull of  $f$  defined by  $h_\lambda f = -e_{\lambda,2}(-e_{\lambda,2}f)$  is given by*

$$-e_{\lambda,2}(-e_{\lambda,2}f) = \left(f + \frac{1}{2\lambda} \|\cdot\|^2\right)^{**} - \frac{1}{2\lambda} \|\cdot\|^2. \tag{54}$$

- (b) *A function  $f : X \rightarrow \overline{\mathbb{R}}$  is a Moreau envelope with index  $\lambda$  and power 2 if and only if  $f - (1/2\lambda)\|\cdot\|^2 \in -\Gamma(X)$ .*

*Proof.* (a) For every  $x \in X$ , we have

$$\begin{aligned}
 e_{\lambda,2}f(x) &= \inf_{y \in X} \left\{ \frac{1}{2\lambda} \|x - y\|^2 + f(y) \right\} \\
 &= \inf_{y \in X} \left\{ \frac{1}{2\lambda} \|x\|^2 + \frac{1}{2\lambda} \|y\|^2 - \frac{1}{\lambda} \langle x, y \rangle + f(y) \right\} \\
 &= -\left(f + \frac{1}{2\lambda} \|\cdot\|^2\right)^* (x/\lambda) + \frac{1}{2\lambda} \|x\|^2,
 \end{aligned}$$

which proves the equality (53). By iterating we deduce that

$$\begin{aligned}
 -e_{\lambda,2}(-e_{\lambda,2}f) &= \left(-e_{\lambda,2}f + \frac{1}{2\lambda} \|\cdot\|^2\right)^* \left(\frac{\cdot}{\lambda}\right) - \frac{1}{2\lambda} \|\cdot\|^2 \\
 &= \left[\left(f + \frac{1}{2\lambda} \|\cdot\|^2\right)^* \left(\frac{\cdot}{\lambda}\right)\right]^* \left(\frac{\cdot}{\lambda}\right) - \frac{1}{2\lambda} \|\cdot\|^2 \\
 &= \left(f + \frac{1}{2\lambda} \|\cdot\|^2\right)^{**} - \frac{1}{2\lambda} \|\cdot\|^2,
 \end{aligned}$$

which proves the equality (54).

(b) Observe that  $f$  is a Moreau envelope with index  $\lambda$  and power 2 if and only if  $-f \in \mathcal{E}^\varphi$  with  $\varphi = -(1/2\lambda)\|\cdot\|^2$ . From the equivalence (7)  $\Leftrightarrow$  (8) and the fact that  $\varphi_- = \varphi$ , this is, in turn, equivalent to  $-f = ((-f)^\varphi)^\varphi$ . Since  $((-f)^\varphi)^\varphi = -e_{\lambda,2}(-e_{\lambda,2}(-f))$  and using the equality (54), we infer that

$f$  is a Moreau envelope with index  $\lambda$  and power 2

$$\begin{aligned}
 &\Downarrow \\
 -f + \frac{1}{2\lambda} \|\cdot\|^2 &= \left(-f + \frac{1}{2\lambda} \|\cdot\|^2\right)^{**} \\
 &\Downarrow \\
 -f + \frac{1}{2\lambda} \|\cdot\|^2 &\in \Gamma(X). \\
 &\Downarrow \\
 f - \frac{1}{2\lambda} \|\cdot\|^2 &\in -\Gamma(X). \quad \square
 \end{aligned}$$

The coupling functional  $(x, y) \mapsto -(1/2\lambda)\|x - y\|^2$  was considered in [7, §5] in the framework of generalized conjugacy. Equalities (53)–(54) were established by Penot and Volle [24, p. 206] and Martinez-Legaz [17, pp. 182–184]. These equalities were also observed in [30, Example 11.26(c)] and [39, Lemma 3.3]. The characterization (b) above has been noted in the aforementioned references, and it amounts to the previous characterization in [7, p. 288] of  $Q^c$ -convex functions with  $c := 1/(2\lambda)$ . It seems that Moreau was the first who provided this characterization (b), see the equivalence (I)  $\Leftrightarrow$  (III) in [20, Proposition 9.b].

### References

1. H. Attouch, G. Buttazzo and G. Michaille, Variational analysis in Sobolev and BV spaces. In *Applications to PDE's and Optimization (MPS/SIAM Series on Optimization 6)*, Society for Industrial and Applied Mathematics (SIAM) (Philadelphia, PA, 2006).
2. A. Auslender and M. Teboulle, *Asymptotic Cones and Functions in Optimization and Variational Inequalities (Springer Monographs in Mathematics)*, Springer (New York, 2003).
3. D. Aze and M. Volle, Various continuity properties of the deconvolution. In *Advances in Optimization (Lectures Notes in Economics and Mathematical Systems 382)*, Springer (1992), 16–30.
4. E. J. Balder, An extension of duality-stability relations to nonconvex optimization problems. *SIAM J. Control Optim.* **15** (1977), 329–343.
5. A. Brønsted and R. T. Rockafellar, On the subdifferentiability of convex functions. *Proc. Amer. Math. Soc.* **16** (1965), 605–611.



6. S. Dolecki, Polarities and generalized extremal convolutions. *J. Convex Anal.* **23** (2016), 603–614.
7. S. Dolecki and S. Kurcyusz, On  $\Phi$ -convexity in extremal problems. *SIAM J. Control Optim.* **16** (1978), 277–300.
8. I. Ekeland and R. Temam, Convex analysis and variational problems. In *SIAM Classics in Applied Mathematics (EkeTem 28)*, Society for Industrial and Applied Mathematics (Philadelphia, PA, 1999).
9. K.-H. Elster and A. Wolf, Recent results on generalized conjugate functions. In *Trends in Mathematical Optimization*, Birkhäuser (Basel, 1988), 67–78.
10. A. S. Granero, M. Jiménez-Sevilla and J. P. Moreno, Intersections of closed balls and geometry of Banach spaces. *Extracta Math.* **19**(1) (2004), 55–92.
11. J.-B. Hiriart-Urruty, A general formula on the conjugate of the difference of functions. *Canad. Math. Bull.* **29** (1986), 482–485.
12. J.-B. Hiriart-Urruty, The deconvolution operation in convex analysis: an introduction. *Cybernet. Systems Anal.* **30** (1994), 555–560.
13. J.-B. Hiriart-Urruty and C. Lemaréchal, Convex analysis and minimization algorithms, I. Fundamentals, II. In *Advanced Theory and Bundle Methods*, Springer (Berlin, 1993).
14. J.-B. Hiriart-Urruty and M.-L. Mazure, Formulations variationnelles de l'addition parallèle et de la soustraction parallèle d'opérateurs semi-définis positifs. *C. R. Acad. Sci. Paris, Sér. I* **302** (1986), 527–530.
15. G. E. Ivanov, Weak convexity of functions and the infimal convolution. *J. Convex Anal.* **23** (2016), 719–732.
16. A. Jourani, L. Thibault and D. Zagrodny, The NSLUC property and Klee envelope. *Math. Ann.* **365**(3–4) (2016), 923–967.
17. J.-E. Martinez-Legaz, Generalized conjugation and related topics. In *'Generalized Convexity and Fractional Programming with Economic Applications'*, *Proc. Pisa. Italy (1988) (Lecture Notes in Economics and Mathematical Systems 345)* (ed. A. Cambini *et al*), Springer (Berlin, 1990), 168–197.
18. J.-E. Martinez-Legaz and J.-P. Penot, Regularization by erasement. *Math. Scand.* **98** (2006), 97–124.
19. S. Mazur, Über schwache Konvergenz in den Räumen  $L^p$ . *Studia Math.* **4** (1933), 128–133.
20. J. J. Moreau, Proximité et dualité dans un espace hilbertien. *Bull. Soc. Math. France* **93** (1965), 273–299.
21. J. J. Moreau, Inf-convolution, sous-additivité, convexité des fonctions numériques. *J. Math. Pures Appl.* **49** (1970), 109–154.
22. J. J. Moreau, Fonctionnelles convexes. In *Collège de France, Paris (1967)*, 2nd edn., Consiglio Nazionale delle Ricerche and Facoltà di Ingegneria Università di Roma 'Tor Vergata' (2003).
23. J.-P. Penot, *Calculus Without Derivatives (Graduate Texts in Mathematics)*, Springer (New York, 2013).
24. J.-P. Penot and M. Volle, On strongly convex and paraconvex dualities. In *'Generalized Convexity and Fractional Programming with Economic Applications'*, *Proc. Pisa. Italy (1988) (Lecture Notes in Economics and Mathematical Systems 345)* (ed. A. Cambini *et al*), Springer (Berlin, 1990), 198–218.
25. E. S. Polovinkin, On strongly convex sets and strongly convex functions. *J. Math. Sci.* **100** (2000), 2633–2681.
26. E. S. Polovinkin and M. V. Balashov, *Elements of Convex and Strongly Convex Analysis*, Fizmatlit (Moscow, 2004) (Russian).
27. B. N. Pshenichnyi, Leçons sur les jeux différentiels. *Cah. I.I.R.I.A.*(4) (1971), 145–226.
28. R. T. Rockafellar, *Convex Analysis*, Princeton University Press (Princeton, NJ, 1970).
29. R. T. Rockafellar, Augmented Lagrange multipliers functions and duality in nonconvex programming. *SIAM J. Control Optim.* **12** (1974), 268–285.
30. R. T. Rockafellar and R. J.-B. Wets, *Variational Analysis*, Springer (Berlin, 1998).
31. A. Rubinov, *Abstract Convexity and Global Optimization*, Kluwer (Dordrecht, 2000).
32. I. Singer, Conjugation operators. In *Selected Topics in Operations Research and Mathematical Economics* (eds G. Hammer and D. Pallaschke), Springer (Berlin, 1984), 80–97.
33. I. Singer, Some Relations between Dualities, Polarities, Coupling Functionals, and Conjugations. *J. Math. Anal. Appl.* **115** (1986), 1–22.
34. I. Singer, *Duality for Nonconvex Approximation and Optimization (CMS Books in Mathematics)*, Springer (New York, 2006).
35. L. Vesely, Affine mappings and convex functions. Examples of convex functions. *Preprint*, available at <http://www.mat.unimi.it/users/libor/AnConvessa/functions.pdf>.
36. J.-P. Vial, Strong and weak convexity of sets and functions. *Math. Oper. Res.* **8** (1983), 231–259.
37. M. Volle, Contributions à la Dualité et à l'Épiconvergence. *Thèse de Doctorat d'État*, Université de Pau et des Pays de l'Adour, 1986.

38. M. Volle, A formula on the subdifferential of the deconvolution of convex functions. *Bull. Aust. Math. Soc.* **47** (1993), 333–340.
39. X. Wang, On Chebyshev functions and Klee functions. *J. Math. Anal. Appl.* **368** (2010), 293–310.

A. Cabot,  
Institut de Mathématiques de Bourgogne,  
UMR 5584,  
CNRS,  
Université Bourgogne Franche-Comté,  
21000 Dijon,  
France  
E-mail: alexandre.cabot@u-bourgogne.fr

A. Jourani,  
Institut de Mathématiques de Bourgogne,  
UMR 5584,  
CNRS,  
Université Bourgogne Franche-Comté,  
21000 Dijon,  
France  
E-mail: abderrahim.jourani@u-bourgogne.fr

L. Thibault,  
Institut Montpellierain Alexander Grothendieck,  
Université de Montpellier,  
Place Eugène Bataillon,  
34095 Montpellier,  
France  
and  
Centro de Modelamiento Matemático  
Universidad de Chile, Chile  
E-mail: lionel.thibault@univ-montp2.fr